

Finite Element Approximation of a Phase Field Model for Surface Diffusion of Voids in a Stressed Solid

John W. Barrett[†] Harald Garcke[‡] Robert Nürnberg[†]

Abstract

We consider a fully practical finite element approximation of the degenerate Cahn–Hilliard equation with elasticity: Find the conserved order parameter, $\theta(x, t) \in [-1, 1]$, and the displacement field, $\underline{u}(x, t) \in \mathbb{R}^2$, such that

$$\begin{aligned} \gamma \frac{\partial \theta}{\partial t} &= \nabla \cdot (b(\theta) \nabla [-\gamma \Delta \theta + \gamma^{-1} \Psi'(\theta) + \frac{1}{2} c'(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) : \underline{\underline{\mathcal{E}}}(\underline{u})]), \\ \nabla \cdot (c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u})) &= \underline{0}, \end{aligned}$$

subject to an initial condition $\theta^0(\cdot) \in [-1, 1]$ on θ and boundary conditions on both equations. Here $\gamma \in \mathbb{R}_{>0}$ is the interfacial parameter, Ψ is a non-smooth double well potential, $\underline{\underline{\mathcal{E}}}$ is the symmetric strain tensor, \mathcal{C} is the possibly anisotropic elasticity tensor, $c(s) := c_0 + \frac{1}{2} (1 - c_0) (1 + s)$ with $c_0(\gamma) \in \mathbb{R}_{>0}$ and $b(s) := 1 - s^2$ is the degenerate diffusional mobility. In addition to showing stability bounds for our approximation; we prove convergence, and hence existence of a solution to this nonlinear degenerate parabolic system in two space dimensions. Finally, some numerical experiments are presented.

Mathematics Subject Classification (2000): 65M60, 65M12, 65M50, 35K55, 35K65, 35K35, 82C26, 74F15

1 Introduction

Integrated circuits contain thin metallic lines (interconnects) that make electrical contact between different components of the device. These lines are passivated with a layer of oxide at large temperatures and during the cooling process large stresses are induced. Also voids nucleate in the interconnect and they migrate and change their shape due to the diffusion of atoms. One of the major failure mechanisms in modern micro-electronic circuits is that voids cut the whole interconnect and cause an open circuit. The understanding of how voids migrate is therefore of great practical interest.

[†]Department of Mathematics, Imperial College, London, SW7 2AZ, U.K.

[‡]NWF I – Mathematik, Universität Regensburg, 93040 Regensburg, Germany

In general diffusion in the bulk of the interconnect is much slower than that on the surface of the void. Therefore we will restrict ourselves to the case where diffusion is restricted to the surface of the void or more precisely to a diffuse layer at the void surface. In this case there are three main driving forces for diffusion: one is resulting from capillary effects and the two others are due to electromigration and stressmigration. To formulate the latter two we need to introduce the electric potential ϕ , the displacement field \underline{u} , the symmetric strain tensor $\underline{\underline{\xi}}(\underline{u}) := \frac{1}{2}(\nabla \underline{u} + (\nabla \underline{u})^T)$ and the elastic energy density $E(\underline{u}) := \frac{1}{2} \underline{\underline{C}} \underline{\underline{\xi}}(\underline{u}) : \underline{\underline{\xi}}(\underline{u})$. Here $\underline{\underline{C}}$ is the possibly anisotropic elasticity tensor, which we assume to be symmetric and positive definite. The product $\underline{\underline{A}} : \underline{\underline{B}}$ of two matrices $\underline{\underline{A}}, \underline{\underline{B}} \in \mathbb{R}^{d \times d}$ is defined as $\sum_{i,j=1}^d \mathcal{A}_{ij} \mathcal{B}_{ij}$. We denote by V the normal velocity of the void surface, $\Gamma(t)$, with the normal pointing into the void and by κ its mean curvature with the sign convention that κ is positive if the interface is curved in the direction of the normal. Then mass conservation gives

$$V = -\nabla_s \cdot \underline{\underline{J}}_s \quad \text{on } \Gamma(t), \quad \text{where} \quad \underline{\underline{J}}_s = -D_s \nabla_s (-\zeta \kappa + E(\underline{u}) + \alpha \phi) \quad (1.1)$$

is the mass flux, $\nabla_s \cdot$ is the surface divergence, ∇_s is the surface gradient, D_s is a constant related to the surface diffusivity and ζ is the surface energy density. Here the first term describes capillary forces, the second describes forces resulting from changes in the elastic energy and the forcing term $\alpha \nabla_s \phi$ is caused by an electric current in the bulk of the material and this force is related to the “electron wind” force. The above equations for the surface motion then have to be coupled to the Laplace equation for the electric potential ϕ , the quasi-static mechanical equilibrium equations for \underline{u} and appropriate boundary conditions. For more details we refer to Xia, Bower, Suo, and Shih (1997), Bower and Craft (1998) and Gungor, Maroudas, and Gray (1998).

Let us briefly discuss the influence of the three terms of the mass flux in (1.1). The first term leads to diffusion of atoms from regions of small mean curvature to regions of high mean curvature. If only capillary effects were present the length/area of the void surface would decrease and the voids would become circular/spherical, see Elliott and Garcke (1997) and Escher, Mayer, and Simonett (1998). The second term leads to diffusion from regions of high elastic energy to regions of smaller elastic energy and the third gives rise to diffusion in the direction opposite to the electric field (this is true if $\alpha < 0$ and this is the case for aluminum, which is mainly used for interconnects). The effect of all three terms can be seen in numerical simulations; see e.g. Bhate, Kumar, and Bower (2000) and Barrett, Nürnberg, and Styles (2004, §5). From these numerical simulations one notices that the topology of the voids can change. Therefore numerical methods that depend on the direct parametrization of the void surface will have difficulties. For an overview on numerical methods for interface motion and their advantages and disadvantages we refer to Elliott (1997).

In this paper we study a finite element approximation of a phase field model for surface diffusion of voids due to capillary effects and stressmigration. We will not include electromigration since a phase field method for surface diffusion in the presence of electromigration (and in the absence of stressmigration) was already analysed in Barrett, Nürnberg, and Styles (2004). A phase field model for electromigration of intergranular

voids, i.e. of voids in solids with different grain orientations, will be discussed in Barrett, Garcke, and Nürnberg (2004a). Furthermore, we will present numerical simulations of the combined effect of surface diffusion, electromigration and stressmigration in a forthcoming paper where we will also discuss applications to epitaxial growth; see Barrett, Garcke, and Nürnberg (2004b).

In a phase field model a diffuse layer is used to describe interfaces or free surfaces. To model surface diffusion by a phase field model we introduce an order parameter θ which (away from a small interfacial layer) attains the value -1 in the void and the value 1 in regions occupied by the metal. In the diffuse interfacial layer θ varies continuously from -1 to 1 . The free energy for the evolution law (1.1) is given by

$$\int_{\Gamma(t)} \varsigma \, ds + \int_{\Omega_+(t)} E(\underline{u}) \, dx ,$$

where the first term is the integral of the surface energy density ς over the void surface and Ω_+ is the region occupied by the metal. In phase field models the surface energy density ς is now replaced by an Ginzburg–Landau free energy density $\varsigma \frac{2}{\pi} [\frac{\gamma}{2} |\nabla\theta|^2 + \gamma^{-1} \Psi(\theta)]$, where γ is a small positive parameter related to the interfacial thickness and Ψ is a free energy density with the two global minima ± 1 . In the above, and throughout, we will use for convenience an obstacle free energy of the form

$$\Psi(s) := \begin{cases} \frac{1}{2}(1 - s^2) & \text{if } s \in [-1, 1], \\ \infty & \text{if } s \notin [-1, 1], \end{cases} \quad (1.2)$$

which restricts the order parameter θ to lie in the interval $[-1, 1]$ and also guarantees that outside a small interfacial layer θ attains the values ± 1 ; see e.g. Blowey and Elliott (1991).

The elastic energy density also has to take the interfacial layer into account and is hence given by

$$E(\theta, \underline{u}) := \frac{1}{2} c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) : \underline{\underline{\mathcal{E}}}(\underline{u}) , \quad (1.3)$$

where c is an interpolation function given by

$$c(s) := c_0 + \frac{1}{2}(1 - c_0)(1 + s) . \quad (1.4)$$

Here $c_0 = c_0(\gamma) \in (0, 1)$ is small and we will assume that $c_0(\gamma) \rightarrow 0$ as $\gamma \rightarrow 0$. Hence, c is affine linear with $c(-1) = c_0 \leq c(s) \leq 1 = c(1)$ for all $s \in [-1, 1]$. Now the total free energy for the phase field model is given by

$$J(\theta, \underline{u}) := \int_{\Omega} [\varsigma \frac{2}{\pi} \{ \frac{\gamma}{2} |\nabla\theta|^2 + \gamma^{-1} \Psi(\theta) \} + E(\theta, \underline{u})] \, dx$$

with the possible addition of an integral over the boundary of Ω , depending on the imposed boundary conditions on \underline{u} .

Now we define the chemical potential, w , via the first variation of J with respect to θ

$$w = \frac{\delta J}{\delta \theta} = [\varsigma \frac{2}{\pi} (-\gamma \Delta\theta + \gamma^{-1} \Psi'(\theta)) + \frac{1}{2} c'(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) : \underline{\underline{\mathcal{E}}}(\underline{u})] \quad (1.5)$$

which is the diffusion potential for θ . The diffusion equation for θ is then given by

$$\gamma \frac{\partial \theta}{\partial t} = \nabla \cdot \left(\frac{8}{\pi} D_s b(\theta) \nabla w \right), \quad (1.6)$$

where

$$b(s) := 1 - s^2 \quad \forall s \in [-1, 1] \quad (1.7)$$

is a degenerate mobility which is zero outside of the interfacial layer. Hence diffusion is restricted to the interfacial layer, which is conceptually close to the idea of surface diffusion where diffusion only takes place on the surface. In fact it was shown by Cahn, Elliott, and Novick-Cohen (1996), using formally matched asymptotic expansions, that (in the absence of elastic effects) the phase field equations as stated above converge, as the interfacial parameter $\gamma \rightarrow 0$, to motion by surface diffusion.

If we include elasticity and require quasi-static equilibrium, i.e.

$$\nabla \cdot (c(\theta) \mathcal{C} \underline{\underline{\boldsymbol{\xi}}(\underline{\mathbf{u}})}) = \underline{\mathbf{0}}, \quad (1.8)$$

we obtain in the limit $\gamma \rightarrow 0$ and $c_0(\gamma) \rightarrow 0$ that the zero level sets of θ converge to a hypersurface $\Gamma(t)$ that evolves according to the law

$$V = D_s \Delta_s \left[-\varsigma \kappa + \frac{1}{2} \mathcal{C} \underline{\underline{\boldsymbol{\xi}}(\underline{\mathbf{u}})} : \underline{\underline{\boldsymbol{\xi}}(\underline{\mathbf{u}})} \right] \quad \text{on } \Gamma(t).$$

This can be shown using formally matched asymptotic expansions when one combines the approaches of Cahn, Elliott, and Novick-Cohen (1996), Leo, Lowengrub, and Jou (1998) and Fried and Gurtin (1994)

The system (1.5)–(1.8) is a degenerate Cahn–Hilliard equation coupled to an elasticity system. If $\mathcal{C} \equiv 0$, then (1.5)–(1.8) collapses to the degenerate Cahn–Hilliard equation without elasticity. Existence of a solution to this fourth order degenerate parabolic equation for θ , as $b(\theta)$ can take on zero values, can be found in Elliott and Garcke (1996). Degenerate parabolic equations of higher order exhibit some new characteristic features which are fundamentally different to those for second order degenerate parabolic equations. The key point is that there is no maximum or comparison principle for parabolic equations of higher order. This drastically complicates the analysis since a lot of results which are known for second order equations are proven with the help of comparison techniques. Related to this is the fact that there is no uniqueness result known for (1.5)–(1.7) with $\mathcal{C} \equiv 0$. Although there is no comparison principle, one of the main features of the system (1.5)–(1.7) is the fact that one can show existence of a solution with $|\theta| \leq 1$ if given initial data $|\theta^0| \leq 1$. This is in contrast to linear parabolic equations of fourth order.

In the case of a constant mobility, i.e. $b(\theta) \equiv 1$, the system (1.5), (1.6) and (1.8) was studied analytically by Garcke (2000), Garcke (2003) and Carrive, Miranville, and Piétrus (2000). For a finite element approximation in this non-degenerate case, see e.g. Garcke, Rumpf, and Weikard (2001) and Garcke and Weikard (2004).

There is very little work on the numerical analysis of degenerate parabolic equations of fourth order: for work on the thin film equation see Barrett, Blowey, and Garcke

(1998), Zhornitskaya and Bertozzi (2000) and Grün and Rumpf (2000), for thin film flows in the presence of surfactants see Barrett, Garcke, and Nürnberg (2003); and for work on degenerate Cahn–Hilliard systems see Barrett, Blowey, and Garcke (1999), Barrett, Blowey, and Garcke (2001) and Barrett and Blowey (2001). In all of these papers, although stability bounds were proved in one and two space dimensions, the main convergence result was restricted to one space dimension. However, recently Grün (2003) has proved convergence in two space dimensions of a finite element approximation to the thin film equation. This approach was extended in (i) Barrett and Nürnberg (2004) and (ii) Barrett, Nürnberg, and Styles (2004) to prove convergence in two space dimensions of a finite element approximation to (i) the thin film equation in the presence of surfactants and repulsive van der Waals forces, and (ii) the phase field approximation of (1.1) in the absence of stressmigration. It is the aim of this paper to propose and prove convergence of a finite element approximation of the degenerate system (1.5)–(1.8), and hence prove existence of a solution to (1.5)–(1.8). Since in the stressmigration case a term that is quadratic in the gradient of \underline{u} — as opposed to linear in ϕ in the electromigration case — appears in the chemical potential w , see (1.5); this makes the convergence analysis in the presence of stressmigration far more complicated than that of electromigration.

Due to a lack of embedding properties, our convergence analysis is restricted to two spatial dimensions (i.e. $d = 2$). For ease of exposition, we will restrict our presentation throughout to this case. However, the phase field approach and the corresponding finite element approximation with the basic energy bound, see (2.80a) below, are easily extended to three spatial dimensions. We adopt the following notation throughout. The trace of a tensor $\underline{\underline{A}}$ is denoted by $\text{Tr}(\underline{\underline{A}}) := \mathcal{A}_{11} + \mathcal{A}_{22}$, and the divergence is defined as $\nabla \cdot \underline{\underline{A}} = (\frac{\partial \mathcal{A}_{11}}{\partial x_1} + \frac{\partial \mathcal{A}_{12}}{\partial x_2}, \frac{\partial \mathcal{A}_{21}}{\partial x_1} + \frac{\partial \mathcal{A}_{22}}{\partial x_2})^T$; see e.g. Brenner and Scott (2002, Chapter 11). We will assume throughout for all $i, j, k, l \in \{1, 2\}$ that

$$(i) \quad \mathcal{C}_{ijkl} = \mathcal{C}_{jikl} = \mathcal{C}_{ijlk} \quad \text{and} \quad (ii) \quad \mathcal{C}_{ijkl} = \mathcal{C}_{klij}. \quad (1.9)$$

Here (i) follows, without loss of generality, from the fact that \mathcal{C} maps symmetric tensors to symmetric tensors; and (ii) follows from the symmetry assumption $\mathcal{C} \underline{\underline{A}} : \underline{\underline{B}} = \underline{\underline{A}} : \mathcal{C} \underline{\underline{B}}$. We assume also throughout that \mathcal{C} is positive definite; that is, there exist constants $m_c, M_c > 0$ such that

$$0 < m_c (\underline{\underline{A}} : \underline{\underline{A}}) \leq \mathcal{C} \underline{\underline{A}} : \underline{\underline{A}} \leq M_c (\underline{\underline{A}} : \underline{\underline{A}}) \quad \forall \underline{\underline{A}} \in \mathbb{R}^{2 \times 2} \setminus \{\underline{\underline{0}}\}. \quad (1.10)$$

If one further assumes cubic symmetry, it holds also that $\mathcal{C}_{1111} = \mathcal{C}_{2222}$ and $\mathcal{C}_{2212} = \mathcal{C}_{1112} = 0$; see e.g. Gurtin (1972). For an isotropic material we obtain that

$$\mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) = 2\mu \underline{\underline{\mathcal{E}}}(\underline{u}) + \lambda \text{Tr}(\underline{\underline{\mathcal{E}}}(\underline{u})) \underline{\underline{\mathcal{I}}}, \quad (1.11)$$

where $\underline{\underline{\mathcal{I}}}$ is the identity tensor, and $\mu \in \mathbb{R}_{>0}$ and $\lambda \in \mathbb{R}_{\geq 0}$ are the Lamé moduli. In what follows, to simplify the presentation, we will set, without loss of generality, the surface diffusivity $D_s = \frac{\pi}{8}$ and the surface energy density $\varsigma = \frac{\pi}{2}$.

In the following we will analyse a finite element approximation of the nonlinear degenerate parabolic system for a given $\gamma \in \mathbb{R}_{>0}$:

(P) Find functions $\theta : \Omega \times [0, T] \rightarrow [-1, 1]$, $w : \Omega \times [0, T] \rightarrow \mathbb{R}$ and $\underline{u} : \Omega \times [0, T] \rightarrow \mathbb{R}^2$ such that

$$\gamma \frac{\partial \theta}{\partial t} = \nabla \cdot (b(\theta) \nabla w) \quad \text{in } \Omega_T, \quad (1.12a)$$

$$w = -\gamma \Delta \theta + \gamma^{-1} \Psi'(\theta) + \frac{1}{2} c'(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) : \underline{\underline{\mathcal{E}}}(\underline{u}) \quad \text{on } \{|\theta| < 1\}, \quad (1.12b)$$

$$\nabla \theta \cdot \underline{\nu} = b(\theta) \nabla w \cdot \underline{\nu} = 0 \quad \text{on } \partial \Omega \times (0, T], \quad (1.12c)$$

$$\theta(x, 0) = \theta^0(x) \in [-1, 1] \quad \forall x \in \Omega, \quad (1.12d)$$

$$\nabla \cdot (c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u})) = \underline{0} \quad \text{in } \Omega_T, \quad c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) \underline{\nu} = \underline{g} \quad \text{on } \partial \Omega \times (0, T]; \quad (1.12e)$$

where Ω is a Lipschitz domain in \mathbb{R}^2 with $\underline{\nu}$ the outward unit normal to its boundary $\partial \Omega$, $T > 0$ is a fixed positive time, and $\Omega_T := \Omega \times (0, T]$. The function $\underline{g} \in \underline{L}^2(\partial \Omega)$ is the given boundary force satisfying the necessary compatibility conditions, $\int_{\partial \Omega} \underline{g} \, ds = \underline{0}$ and $\int_{\partial \Omega} \underline{g} \cdot (x_2, -x_1)^T \, ds = 0$, for the existence of a solution \underline{u} to (1.12e). For simplicity, we will consider

$$\underline{g} = \underline{\underline{S}} \underline{\nu} = \mathcal{C} \underline{\underline{S}}^* \underline{\nu}; \quad (1.13)$$

where $\underline{\underline{S}} \in \mathbb{R}^{2 \times 2}$ is a symmetric tensor and $\underline{\underline{S}}^* := \mathcal{C}^{-1} \underline{\underline{S}}$. Alternatively, one could prescribe displacement boundary conditions, $\underline{u} = \underline{f}$, on $\partial \Omega$ or on parts thereof.

We should note that the solution \underline{u} to (1.12e) is not unique. This is simply because

$$\underline{\underline{\mathcal{E}}}(\underline{v}) = \underline{0} \quad \forall \underline{v} \in \underline{\underline{\mathbf{RM}}}, \quad (1.14)$$

where $\underline{\underline{\mathbf{RM}}}$ is the space of rigid motions and characterized by

$$\underline{\underline{\mathbf{RM}}} := \{\underline{v} \in \underline{H}^1(\Omega) : \underline{v} = \underline{p} + q(x_2, -x_1)^T \quad \underline{p} \in \mathbb{R}^2, q \in \mathbb{R}\}.$$

Hence one can impose uniqueness for (1.12e) by seeking \underline{u} such that $\int_{\Omega} \underline{u} \cdot \underline{\nu} \, dx = 0$ for all $\underline{v} \in \underline{\underline{\mathbf{RM}}}$; see our definition of $\widehat{\underline{V}}_p$ in (1.20) below.

The basic ingredients of our approach are some key energy estimates. First, we relate G to b by the identity

$$b(s) G''(s) = 1. \quad (1.15)$$

Knowing b , recall (1.7), the above identity determines G up to a linear term. Furthermore we have that G is convex. One can then derive formally the following energy estimates for (P). Testing (1.12a) with w and (1.12b) with $\frac{\partial \theta}{\partial t}$, combining and noting (1.12c,e) and (1.3) yields that

$$\frac{d}{dt} \left\{ \int_{\Omega} \left[\frac{1}{2} \gamma |\nabla \theta|^2 + \gamma^{-1} \Psi(\theta) + E(\theta, \underline{u}) \right] dx - \int_{\partial \Omega} \underline{g} \cdot \underline{u} \, ds \right\} + \gamma^{-1} \int_{\Omega} b(\theta) |\nabla w|^2 \, dx \leq 0. \quad (1.16)$$

Testing (1.12a) with $G'(\theta)$ and (1.12b) with $-\Delta \theta$, combining and noting (1.15), (1.2) (1.10) and (1.4) yields that

$$\gamma^2 \frac{d}{dt} \int_{\Omega} G(\theta) \, dx + \frac{1}{2} \gamma^2 \int_{\Omega} |\Delta \theta|^2 \, dx \leq \int_{\Omega} |\nabla \theta|^2 \, dx + \frac{1}{32} M_{\mathcal{C}}^2 \int_{\Omega} |\underline{\underline{\mathcal{E}}}(\underline{u})|^4 \, dx. \quad (1.17)$$

In order to bound $\Delta\theta$ in $L^2(\Omega_T)$, one needs to bound $\nabla \underline{u}$ in $L^4(\Omega_T)$. This is the key difficulty when including the elastic effects. This is achieved by using an $L^\infty(0, T; \underline{W}^{1,p}(\Omega))$, $p > 2$, bound for \underline{u} solving (1.12e) which does not depend on the choice of $\theta \in L^\infty(\Omega_T)$, see Garcke (2000), Garcke (2004) and Lemma 1.1 together with Remark 2.2 below.

It is the goal of this paper to derive a finite element approximation of (P) that is consistent with the energy estimates (1.16) and (1.17). In order to derive a discrete analogue of the energy estimate (1.17), we adapt a technique introduced in Zhornitskaya and Bertozzi (2000) and Grün and Rumpf (2000) for deriving a discrete entropy bound for the thin film equation; see also Barrett, Nürnberg, and Styles (2004). However, the key difficulty here in proving convergence of our finite element approximation, and hence existence of a solution to (P), is the finite element analogue of the crucial $\underline{W}^{1,p}(\Omega)$, $p > 2$, bound for \underline{u} ; see Lemma 2.3 below.

This paper is organised as follows. In §2 we formulate a fully practical finite element approximation of the degenerate system (P) and derive discrete analogues of the energy estimates (1.16) and (1.17). In §3 we prove convergence, and hence existence of a solution to the system (P) in two space dimensions. Finally, in §4 we present some numerical experiments.

Notation and Auxiliary Results

Let $D \subset \mathbb{R}^d$, $d = 1$ or 2 , with a Lipschitz boundary ∂D if $d = 2$. We adopt the standard notation for Sobolev spaces, denoting the norm of $W^{m,q}(D)$ ($m \in \mathbb{N}$, $q \in [1, \infty]$) by $\|\cdot\|_{m,q,D}$ and the semi-norm by $|\cdot|_{m,q,D}$. We extend these norms and semi-norms in the natural way to the corresponding spaces of vector and matrix valued functions, which will be denoted by e.g. $\underline{W}^{m,q}(D)$. For $q = 2$, $W^{m,2}(D)$ will be denoted by $H^m(D)$ with the associated norm and semi-norm written as, respectively, $\|\cdot\|_{m,D}$ and $|\cdot|_{m,D}$. For notational convenience, we drop the domain subscript on the above norms and semi-norms in the case $D \equiv \Omega$. Throughout (\cdot, \cdot) denotes the standard L^2 inner product over Ω . In addition we define

$$\mathcal{f}\eta := \frac{1}{\underline{m}(\Omega)} (\eta, 1) \quad \forall \eta \in L^1(\Omega).$$

For later purposes, we recall the following well-known Sobolev interpolation result, e.g. see Adams and Fournier (1977): Let $q \in (1, \infty)$, $r \in [q, \infty)$ if $q \geq 2$ and $r \in [q, \frac{2q}{2-q}]$ if $q \in (1, 2)$; and $\mu := \frac{2}{q} - \frac{2}{r}$. Then the following inequality holds

$$|z|_{0,r} \leq C |z|_{0,q}^{1-\mu} \|z\|_{1,q}^\mu \quad \forall z \in W^{1,q}(\Omega). \quad (1.18)$$

We recall also the following compactness results. Let X , Y and Z be Banach spaces with a compact embedding $X \hookrightarrow Y$ and a continuous embedding $Y \hookrightarrow Z$. Then the embeddings

$$\{ \eta \in L^2(0, T; X) : \frac{\partial \eta}{\partial t} \in L^2(0, T; Z) \} \hookrightarrow L^2(0, T; Y) \quad (1.19a)$$

$$\text{and} \quad \{ \eta \in L^\infty(0, T; X) : \frac{\partial \eta}{\partial t} \in L^2(0, T; Z) \} \hookrightarrow C([0, T]; Y) \quad (1.19b)$$

are compact and a generalised version of (1.19a), where the time derivative is replaced by a time translation, holds. That is, any bounded and closed subset E of $L^2(0, T; X)$ with

$$\lim_{\vartheta \rightarrow 0} \left\{ \sup_{\eta \in E} \|\eta(\cdot, \cdot + \vartheta) - \eta(\cdot, \cdot)\|_{L^2(0, T-\vartheta; Z)} \right\} = 0 \quad (1.19c)$$

is compact in $L^2(0, T; Y)$, see Simon (1987).

For $p \in [1, \infty]$, we introduce also

$$\widehat{V}_p := \{\underline{\eta} \in \underline{W}^{1,p}(\Omega) : (\underline{\eta}, \underline{v}) = 0 \quad \forall \underline{v} \in \underline{RM}\} \quad (1.20)$$

and define $\widehat{H}^1(\Omega) := \widehat{V}_2$. We recall the following version of Korn's inequality

$$\|\underline{\eta}\|_{1,p} \leq C |\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,p} \quad \forall \underline{\eta} \in \widehat{V}_p, \quad p \in (1, \infty); \quad (1.21)$$

see e.g. Nečas and Hlaváček (1981, p79) for the case $p = 2$, or Mosolov and Mjasnikov (1971) for general $p \in (1, \infty)$. Moreover, the following lemma holds.

LEMMA. 1.1 *There exists $\delta \in \mathbb{R}_{>0}$ such that for all $p \in [\frac{2+\delta}{1+\delta}, 2+\delta]$ there is a $\beta(p) \geq 1$ satisfying*

$$|\underline{\underline{\mathcal{E}}}(\underline{z})|_{0,p} \leq \beta(p) \sup_{\underline{0} \neq \underline{\eta} \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(\underline{z}), \underline{\underline{\mathcal{E}}}(\underline{\eta}))}{|\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,q}} \quad \forall \underline{z} \in \widehat{V}_p, \quad (1.22)$$

where $\frac{1}{p} + \frac{1}{q} = 1$. Moreover β is continuous on the interval $[\frac{2+\delta}{1+\delta}, 2+\delta]$ and $\beta(p) \rightarrow \beta(2) = 1$ as $p \rightarrow 2$.

Proof. Let $[L^p(\Omega)]_{sym}^{2 \times 2} := \{\underline{\underline{\mathcal{F}}} \in [L^p(\Omega)]^{2 \times 2} : \underline{\underline{\mathcal{F}}}$ is symmetric $\}$. For $\underline{z} \in \widehat{V}_p$ we define $\mathcal{S}(\underline{\underline{\mathcal{F}}}) := (\underline{\underline{\mathcal{E}}}(\underline{z}), \underline{\underline{\mathcal{F}}})$ for all $\underline{\underline{\mathcal{F}}} = \underline{\underline{\mathcal{E}}}(\underline{\eta})$ with $\underline{\eta} \in \widehat{V}_q$. \mathcal{S} is a continuous linear functional on a closed subspace of $[L^q(\Omega)]_{sym}^{2 \times 2}$ with norm

$$\sup_{\underline{0} \neq \underline{\eta} \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(\underline{z}), \underline{\underline{\mathcal{E}}}(\underline{\eta}))}{|\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,q}}.$$

The Hahn–Banach theorem and the fact that $([L^q(\Omega)]_{sym}^{2 \times 2})' \cong [L^p(\Omega)]_{sym}^{2 \times 2}$ imply the existence of a $\underline{\underline{\mathcal{G}}}_z \in [L^p(\Omega)]_{sym}^{2 \times 2}$ such that

$$(\underline{\underline{\mathcal{E}}}(\underline{z}), \underline{\underline{\mathcal{F}}}) = (\underline{\underline{\mathcal{G}}}_z, \underline{\underline{\mathcal{F}}}) \quad \forall \underline{\underline{\mathcal{F}}} \in [L^q(\Omega)]_{sym}^{2 \times 2} \quad \text{and} \quad |\underline{\underline{\mathcal{G}}}_z|_{0,p} = \sup_{\underline{0} \neq \underline{\eta} \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(\underline{z}), \underline{\underline{\mathcal{E}}}(\underline{\eta}))}{|\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,q}}. \quad (1.23)$$

Let $\mathcal{Q} : [L^p(\Omega)]_{sym}^{2 \times 2} \rightarrow [L^p(\Omega)]_{sym}^{2 \times 2}$ be the linear operator such that $\mathcal{Q} \underline{\underline{\mathcal{F}}} = \underline{\underline{\mathcal{E}}}(f_{\underline{\mathcal{F}}})$, where $f_{\underline{\mathcal{F}}} \in \widehat{V}_p$ is such that

$$(\underline{\underline{\mathcal{E}}}(f_{\underline{\mathcal{F}}}), \underline{\underline{\mathcal{E}}}(\underline{\eta})) = (\underline{\underline{\mathcal{F}}}, \underline{\underline{\mathcal{E}}}(\underline{\eta})) \quad \forall \underline{\eta} \in \widehat{V}_q. \quad (1.24)$$

We need to show that \mathcal{Q} is well-defined and compute the operator norm $\|\mathcal{Q}\|_p$ of \mathcal{Q} . The well-posedness of \mathcal{Q} for $p = 2$ follows from (1.21) and the Lax–Milgram theorem; and in addition, $\|\mathcal{Q}\|_2 = 1$. Moreover, regularity theory implies that there exists a $\delta > 0$ such that for all $p \in [2, 2 + \delta]$ if $\underline{\underline{\mathcal{F}}} \in [L^p(\Omega)]_{sym}^{2 \times 2}$ then it holds that

$$|\mathcal{Q}\underline{\underline{\mathcal{F}}}|_{0,p} = |\underline{\underline{\mathcal{E}}}(f_{\underline{\underline{\mathcal{F}}}})|_{0,p} \leq C(p) [|\nabla f_{\underline{\underline{\mathcal{F}}}}|_{0,2} + |\underline{\underline{\mathcal{F}}}|_{0,p}] \leq C(p) [|\underline{\underline{\mathcal{F}}}|_{0,2} + |\underline{\underline{\mathcal{F}}}|_{0,p}] \leq C(p) |\underline{\underline{\mathcal{F}}}|_{0,p}.$$

The first inequality in the above can be shown for example with the help of a method introduced by Giaquinta and Modica (1979), who proved local L^p -estimates for gradients of solutions to elliptic systems. In Garcke (2000) and Garcke (2004) this method has been applied to obtain global L^p -estimates for gradients of solutions to elasticity systems on Lipschitz domains. The above shows that \mathcal{Q} is a bounded linear operator for $p \in [2, 2 + \delta]$ and that $\|\mathcal{Q}\|_p \leq C(p)$.

We now want to show that \mathcal{Q} is also a linear continuous operator on $[L^q(\Omega)]_{sym}^{2 \times 2}$, where q is such that $\frac{1}{p} + \frac{1}{q} = 1$ for a $p \in [2, 2 + \delta]$. To do so, we approximate $\underline{\underline{\mathcal{F}}} \in [L^q(\Omega)]_{sym}^{2 \times 2}$ by $\underline{\underline{\mathcal{F}}}_k \in [L^2(\Omega)]_{sym}^{2 \times 2}$ such that $|\underline{\underline{\mathcal{F}}} - \underline{\underline{\mathcal{F}}}_k|_{0,q} \rightarrow 0$ as $k \rightarrow \infty$. As $\widehat{V}_p \subset \widehat{V}_2$ it then follows that

$$(\mathcal{Q}\underline{\underline{\mathcal{F}}}_k, \underline{\underline{\mathcal{H}}}) = (\mathcal{Q}\underline{\underline{\mathcal{F}}}_k, \mathcal{Q}\underline{\underline{\mathcal{H}}}) = (\underline{\underline{\mathcal{F}}}_k, \mathcal{Q}\underline{\underline{\mathcal{H}}}) \quad \forall \underline{\underline{\mathcal{H}}} \in [L^p(\Omega)]_{sym}^{2 \times 2}.$$

Hence we obtain that

$$|(\mathcal{Q}\underline{\underline{\mathcal{F}}}_k, \underline{\underline{\mathcal{H}}})| \leq \|\mathcal{Q}\|_p |\underline{\underline{\mathcal{H}}}|_{0,p} |\underline{\underline{\mathcal{F}}}_k|_{0,q} \implies |\mathcal{Q}\underline{\underline{\mathcal{F}}}_k|_{0,q} \leq \|\mathcal{Q}\|_p |\underline{\underline{\mathcal{F}}}_k|_{0,q}.$$

Taking the weak limit of $f_{\underline{\underline{\mathcal{F}}}_k}$ in \widehat{V}_q , where $\underline{\underline{\mathcal{E}}}(f_{\underline{\underline{\mathcal{F}}}_k}) = \mathcal{Q}\underline{\underline{\mathcal{F}}}_k$, we obtain that $(\mathcal{Q}\underline{\underline{\mathcal{F}}}, \underline{\underline{\mathcal{H}}}) = (\underline{\underline{\mathcal{F}}}, \mathcal{Q}\underline{\underline{\mathcal{H}}})$ for all $\underline{\underline{\mathcal{F}}} \in [L^q(\Omega)]_{sym}^{2 \times 2}$ and $\underline{\underline{\mathcal{H}}} \in [L^p(\Omega)]_{sym}^{2 \times 2}$. Hence \mathcal{Q} defined on $[L^q(\Omega)]_{sym}^{2 \times 2}$ is the dual operator to \mathcal{Q} defined on $[L^p(\Omega)]_{sym}^{2 \times 2}$ and therefore $\|\mathcal{Q}\|_p = \|\mathcal{Q}\|_q$.

The Riesz–Thorin theorem, see Bergh and Löfström (1976), then implies that $\|\mathcal{Q}\|_p \leq \|\mathcal{Q}\|_s^{1-\alpha} \|\mathcal{Q}\|_r^\alpha$ for all $\frac{2+\delta}{1+\delta} \leq s \leq p \leq r \leq 2 + \delta$ such that $\frac{1}{p} = (1-\alpha)\frac{1}{s} + \alpha\frac{1}{r}$ and $\alpha \in [0, 1]$. It follows that $\log \|\mathcal{Q}\|_p$ is a convex function of $\frac{1}{p}$ and therefore $\|\mathcal{Q}\|_p$ is a continuous function of p with $\|\mathcal{Q}\|_2 = 1$. Finally, it follows from (1.23) and (1.24) that $\underline{\underline{\mathcal{E}}}(\underline{\underline{z}}) = \mathcal{Q}\underline{\underline{\mathcal{G}}}_z$ and hence $|\underline{\underline{\mathcal{E}}}(\underline{\underline{z}})|_{0,p} \leq \|\mathcal{Q}\|_p |\underline{\underline{\mathcal{G}}}_z|_{0,p}$. Therefore the desired result (1.22) follows from (1.23) with $\beta(p) = \|\mathcal{Q}\|_p$. \square

We note also for future reference the generalised Young’s inequality

$$r s \leq \frac{1}{p} (\alpha r)^p + \frac{1}{q} (\alpha^{-1} s)^q \quad \forall r, s \in \mathbb{R}, \alpha \in \mathbb{R}_{>0}, p \in (1, \infty) \text{ with } \frac{1}{p} + \frac{1}{q} = 1. \quad (1.25)$$

Throughout C denotes a generic constant independent of h , τ and ε ; the mesh and temporal discretization parameters and the regularization parameter. In addition $C(a_1, \dots, a_I)$ denotes a constant depending on the arguments $\{a_i\}_{i=1}^I$. Finally, $\cdot^{(\star)}$ denotes an expression with or without the superscript \star .

2 Finite Element Approximation

We consider the finite element approximation of (P) under the following assumptions on the mesh:

- (A) Let Ω be a convex polygonal domain. Let $\{\mathcal{T}^h\}_{h>0}$ be a quasi-uniform family of partitionings of Ω into disjoint open simplices σ with $h_\sigma := \text{diam}(\sigma)$ and $h := \max_{\sigma \in \mathcal{T}^h} h_\sigma$, so that $\overline{\Omega} = \cup_{\sigma \in \mathcal{T}^h} \overline{\sigma}$. In addition, it is assumed that all simplices $\sigma \in \mathcal{T}^h$ are right-angled.

We note that the right-angled simplices assumption is not such a severe constraint for appropriate domains Ω , as there exist adaptive finite element codes that satisfy this requirement, see e.g. Schmidt and Siebert (2001).

Associated with \mathcal{T}^h is the finite element space

$$S^h := \{\chi \in C(\overline{\Omega}) : \chi|_\sigma \text{ is linear } \forall \sigma \in \mathcal{T}^h\} \subset H^1(\Omega).$$

We introduce also $\underline{S}^h := [S^h]^2$, $\widehat{S}^h := \underline{S}^h \cap \widehat{H}^1(\Omega)$ and

$$K^h := \{\chi \in S^h : |\chi| \leq 1 \text{ in } \Omega\} \subset K := \{\eta \in H^1(\Omega) : |\eta| \leq 1 \text{ a.e. in } \Omega\}.$$

Let J be the set of nodes of \mathcal{T}^h and $\{p_j\}_{j \in J}$ the coordinates of these nodes. Let $\{\chi_j\}_{j \in J}$ be the standard basis functions for S^h ; that is $\chi_j \in K^h$ and $\chi_j(p_i) = \delta_{ij}$ for all $i, j \in J$. The right angle constraint on the partitioning is required for our approximation of $b(\cdot)$, see (2.9a,b) below, but one consequence is that

$$\int_\sigma \nabla \chi_i \cdot \nabla \chi_j \, dx \leq 0 \quad i \neq j, \quad \forall \sigma \in \mathcal{T}^h. \quad (2.1)$$

We introduce $\pi^h : C(\overline{\Omega}) \rightarrow S^h$, the interpolation operator, such that $(\pi^h \eta)(p_j) = \eta(p_j)$ for all $j \in J$. A discrete semi-inner product on $C(\overline{\Omega})$ is then defined by

$$(\eta_1, \eta_2)^h := \int_\Omega \pi^h(\eta_1(x) \eta_2(x)) \, dx = \sum_{j \in J} m_j \eta_1(p_j) \eta_2(p_j), \quad (2.2)$$

where $m_j := (1, \chi_j) > 0$. The induced discrete semi-norm is then

$$|\eta|_h := [(\eta, \eta)^h]^{\frac{1}{2}} = \left(\int_\Omega \pi^h[\eta^2] \, dx \right)^{\frac{1}{2}} \quad \forall \eta \in C(\overline{\Omega}). \quad (2.3)$$

Both (2.2) and (2.3) are naturally extended to vector and matrix valued functions. We introduce also the L^2 projection $Q^h : L^2(\Omega) \rightarrow S^h$ defined by

$$(Q^h \eta, \chi)^h = (\eta, \chi) \quad \forall \chi \in S^h. \quad (2.4)$$

On recalling (1.7) and (1.15), we then define a function G such that $b(\theta) \nabla[G'(\theta)] = \nabla\theta$; that is,

$$G''(s) = \frac{1}{b(s)} = \frac{1}{1-s^2}. \quad (2.5)$$

We take

$$G(s) = \frac{1}{2}[F(s) + F(-s)], \quad \text{where} \quad F(s) := (1+s) \log(1+s) + (1-s). \quad (2.6)$$

As in Barrett, Nürnberg, and Styles (2004), for computational purposes we replace $F \in C^\infty(-1, \infty)$, $G \in C^\infty(-1, 1)$ for any $\varepsilon \in (0, 1)$ by the regularized functions $F_\varepsilon, G_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$F_\varepsilon(s) := \begin{cases} F(\varepsilon - 1) + (s - \varepsilon + 1) F'(\varepsilon - 1) + \frac{(s - \varepsilon + 1)^2}{2} F''(\varepsilon - 1) & s \leq \varepsilon - 1 \\ F(s) & s \geq \varepsilon - 1 \end{cases}$$

and $G_\varepsilon(s) := \frac{1}{2}[F_\varepsilon(s) + F_\varepsilon(-s)]. \quad (2.7)$

Hence $F_\varepsilon, G_\varepsilon \in C^{2,1}(\mathbb{R})$ with the first two derivatives of F_ε given by

$$F'_\varepsilon(s) := \begin{cases} F'(\varepsilon - 1) + (s - \varepsilon + 1) F''(\varepsilon - 1) & s \leq \varepsilon - 1 \\ F'(s) & s \geq \varepsilon - 1 \end{cases}$$

and $F''_\varepsilon(s) := \begin{cases} F''(\varepsilon - 1) & s \leq \varepsilon - 1 \\ F''(s) & s \geq \varepsilon - 1, \end{cases}$

respectively. We note for later purposes that for all $s \in [-1, 1]$

$$\frac{1}{2} \leq F''_\varepsilon(s) \leq \varepsilon^{-1}, \quad \frac{1}{2} F''_\varepsilon(s) \leq G''_\varepsilon(s) \leq [\varepsilon(2 - \varepsilon)]^{-1} \leq \varepsilon^{-1}. \quad (2.8)$$

Similarly to the approach in Zhornitskaya and Bertozzi (2000) and Grün and Rumpf (2000), we introduce $\Xi_\varepsilon : S^h \rightarrow [L^\infty(\Omega)]^{2 \times 2}$ approximating $b(\cdot) \mathcal{I}$, where $\mathcal{I} \in \mathbb{R}^{2 \times 2}$ is the identity matrix, such that for all $z^h \in S^h$ and *a.e.* in Ω

$$\Xi_\varepsilon(z^h) \text{ is symmetric and positive semi-definite,} \quad (2.9a)$$

$$\Xi_\varepsilon(z^h) \nabla \pi^h[G'_\varepsilon(z^h)] = \nabla z^h. \quad (2.9b)$$

We now give the construction of Ξ_ε . Let $\{e_i\}_{i=1}^2$ be the orthonormal vectors in \mathbb{R}^2 , such that the j^{th} component of e_i is δ_{ij} , $i, j = 1 \rightarrow 2$. Given non-zero constants α_i , $i = 1 \rightarrow 2$; let $\widehat{\sigma}(\{\alpha_i\}_{i=1}^2)$ be the reference open simplex in \mathbb{R}^2 with vertices $\{\widehat{p}_i\}_{i=0}^2$, where \widehat{p}_0 is the origin and $\widehat{p}_i = \alpha_i e_i$, $i = 1 \rightarrow 2$. Given a $\sigma \in \mathcal{T}^h$ with vertices $\{p_{j_i}\}_{i=0}^2$, such that p_{j_0} is the right-angled vertex, then there exists a rotation matrix R_σ and non-zero constants $\{\alpha_i\}_{i=1}^2$ such that the mapping $\mathcal{R}_\sigma : \widehat{x} \in \mathbb{R}^2 \rightarrow p_{j_0} + R_\sigma \widehat{x} \in \mathbb{R}^2$ maps the vertex \widehat{p}_i to p_{j_i} , $i = 0 \rightarrow 2$, and hence $\widehat{\sigma} \equiv \widehat{\sigma}(\{\alpha_i\}_{i=1}^2)$ to σ . For any $z^h \in S^h$, we then set

$$\Xi_\varepsilon(z^h)|_\sigma := R_\sigma \widehat{\Xi}_\varepsilon(\widehat{z}^h)|_{\widehat{\sigma}} R_\sigma^T, \quad (2.10)$$

where $\widehat{z}^h(\widehat{x}) \equiv z^h(\mathcal{R}_\sigma \widehat{x})$ for all $\widehat{x} \in \widehat{\sigma}$ and $\widehat{\Xi}_\varepsilon(\widehat{z}^h)|_{\widehat{\sigma}}$ is the 2×2 diagonal matrix with diagonal entries, $k = 1 \rightarrow 2$,

$$[\widehat{\Xi}_\varepsilon(\widehat{z}^h)|_{\widehat{\sigma}}]_{kk} := \begin{cases} \frac{\widehat{z}^h(\widehat{p}_k) - \widehat{z}^h(\widehat{p}_0)}{G'_\varepsilon(\widehat{z}^h(\widehat{p}_k)) - G'_\varepsilon(\widehat{z}^h(\widehat{p}_0))} \equiv \frac{z^h(p_{j_k}) - z^h(p_{j_0})}{G'_\varepsilon(z^h(p_{j_k})) - G'_\varepsilon(z^h(p_{j_0}))} & \text{if } z^h(p_{j_k}) \neq z^h(p_{j_0}), \\ \frac{1}{G''_\varepsilon(\widehat{z}^h(\widehat{p}_0))} \equiv \frac{1}{G''_\varepsilon(z^h(p_{j_0}))} & \text{if } z^h(p_{j_k}) = z^h(p_{j_0}). \end{cases} \quad (2.11)$$

As $R_\sigma^T \equiv R_\sigma^{-1}$, $\nabla z^h \equiv R_\sigma \widehat{\nabla} \widehat{z}^h$, where $x \equiv (x_1, x_2)^T$, $\nabla \equiv (\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2})^T$, $\widehat{x} \equiv (\widehat{x}_1, \widehat{x}_2)^T$ and $\widehat{\nabla} \equiv (\frac{\partial}{\partial \widehat{x}_1}, \frac{\partial}{\partial \widehat{x}_2})^T$, it easily follows that $\Xi_\varepsilon(z^h)$ constructed in (2.10) and (2.11) satisfies (2.9a,b). We note that it is this construction that requires the right angle constraint on the partitioning \mathcal{T}^h .

In addition to \mathcal{T}^h , let $0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$ be a partitioning of $[0, T]$ into possibly variable time steps $\tau_n := t_n - t_{n-1}$, $n = 1 \rightarrow N$. We set $\tau := \max_{n=1 \rightarrow N} \tau_n$. For any given $\varepsilon \in (0, 1)$, we then consider the following fully practical finite element approximation of (P):

($\mathbf{P}_\varepsilon^{h,\tau}$) For $n \geq 1$ find $\{\underline{U}_\varepsilon^n, \Theta_\varepsilon^n, W_\varepsilon^n\} \in \widehat{\underline{S}}^h \times K^h \times S^h$ such that

$$(c(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\underline{\chi})) = \int_{\partial\Omega} \underline{g} \cdot \underline{\chi} \, ds \quad \forall \underline{\chi} \in \underline{S}^h, \quad (2.12a)$$

$$\gamma \left(\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n}, \chi \right)^h + (\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla \chi) = 0 \quad \forall \chi \in S^h, \quad (2.12b)$$

$$\begin{aligned} \gamma (\nabla \Theta_\varepsilon^n, \nabla [\chi - \Theta_\varepsilon^n]) &\geq (W_\varepsilon^n + \gamma^{-1} \Theta_\varepsilon^{n-1}, \chi - \Theta_\varepsilon^n)^h \\ &\quad - \frac{1}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \chi - \Theta_\varepsilon^n) \quad \forall \chi \in K^h, \end{aligned} \quad (2.12c)$$

where $\Theta_\varepsilon^0 \in K^h$ is an approximation of $\theta^0 \in K$, e.g. $\Theta_\varepsilon^0 \equiv Q^h \theta^0$, or $\Theta_\varepsilon^0 \equiv \pi^h \theta^0$ if $\theta^0 \in C(\overline{\Omega})$.

REMARK. 2.1 We note that in the case $\mathcal{C} \equiv 0$, (2.12b,c) collapses to an approximation of the degenerate Cahn–Hilliard equation, (1.12a–c) with $\mathcal{C} \equiv 0$. This is the same as the approximation in Barrett, Nürnberg, and Styles (2004) in the absence of an electric field. Note that as c' is constant, the dependence on Θ_ε^{n-1} in (2.12c) is superfluous.

Below we recall some well-known results concerning S^h for any $\sigma \in \mathcal{T}^h$, $\chi, z^h \in S^h$, $m \in \{0, 1\}$, $p \in [1, \infty]$, $q \in [2, \infty]$ and $r \in (2, \infty]$:

$$|\chi|_{1,\sigma} \leq C h_\sigma^{-1} |\chi|_{0,\sigma}; \quad (2.13)$$

$$|\chi|_{m,s,\sigma} \leq C h_\sigma^{-2(\frac{1}{p}-\frac{1}{s})} |\chi|_{m,p,\sigma} \quad \text{for any } s \in [p, \infty]; \quad (2.14)$$

$$|(I - \pi^h)\eta|_{m,q,\sigma} \leq C h^{1+\frac{2}{q}-m} |\eta|_{2,\sigma} \quad \forall \eta \in H^2(\sigma); \quad (2.15)$$

$$|(I - \pi^h)\eta|_{m,r,\sigma} \leq C h^{1-m} |\eta|_{1,r,\sigma} \quad \forall \eta \in W^{1,r}(\sigma); \quad (2.16)$$

$$\int_\sigma \chi^2 \, dx \leq \int_\sigma \pi^h[\chi^2] \, dx \leq 4 \int_\sigma \chi^2 \, dx; \quad (2.17)$$

$$|(\chi, z^h) - (\chi, z^h)^h| \leq |(I - \pi^h)(\chi z^h)|_{0,1} \leq C h^{1+m} |\chi|_m |z^h|_1. \quad (2.18)$$

Finally, as we have a quasi-uniform family of partitionings, it holds that

$$|(I - Q^h)\eta|_m \leq C h^{1-m} |\eta|_1 \quad \forall \eta \in H^1(\Omega). \quad (2.19)$$

It is convenient to introduce the “inverse Laplacian” operator $\mathcal{G} : Y \rightarrow Z$ such that

$$(\nabla[\mathcal{G}z], \nabla\eta) = (z, \eta) \quad \forall \eta \in H^1(\Omega), \quad (2.20)$$

where $Y := \{z \in (H^1(\Omega))' : \langle z, 1 \rangle = 0\}$ and $Z := \{z \in H^1(\Omega) : (z, 1) = 0\}$. Here and throughout $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $(H^1(\Omega))'$ and $H^1(\Omega)$, and its extension to the corresponding spaces of vector valued functions. The well-posedness of \mathcal{G} follows from the generalised Lax–Milgram theorem and the Poincaré inequality

$$|\eta|_0 \leq C (|\eta|_1 + |(\eta, 1)|) \quad \forall \eta \in H^1(\Omega). \quad (2.21)$$

As Ω is convex polygonal, we recall the well-known regularity result

$$\|\mathcal{G}z\|_2 \leq C|z|_0 \quad \forall z \in L^2(\Omega) \cap Y. \quad (2.22)$$

We define $Z^h := \{z^h \in S^h : (z^h, 1) = 0\} \subset Y^h := \{z \in C(\bar{\Omega}) : (z, 1)^h = 0\} \subset Y$. Then, similarly to (2.20), we introduce $\mathcal{G}^h : Y^h \rightarrow Z^h$ such that

$$(\nabla[\mathcal{G}^h z^h], \nabla\chi) = (z^h, \chi)^h \quad \forall \chi \in S^h. \quad (2.23)$$

It is easily established from (2.20), (2.23), $\{\mathcal{T}^h\}_{h>0}$ being a regular partitioning, (2.22) and (2.18) that

$$\|(\mathcal{G} - \mathcal{G}^h)z^h\|_1 \leq Ch|z^h|_0 \quad \forall z^h \in S^h. \quad (2.24)$$

We introduce the “discrete Laplacian” operator $\Delta^h : S^h \rightarrow Z^h$ such that

$$(\Delta^h z^h, \chi)^h = -(\nabla z^h, \nabla\chi) \quad \forall \chi \in S^h. \quad (2.25)$$

It follows from (2.2), (2.17), (2.25), (2.14) and the quasi-uniformity assumption on \mathcal{T}^h that

$$\begin{aligned} |\Delta^h z^h|_0^2 &\leq |\Delta^h z^h|_h^2 = -(\nabla z^h, \nabla(\Delta^h z^h)) \leq |z^h|_1 |\Delta^h z^h|_1 \\ &\leq Ch^{-1} |z^h|_1 |\Delta^h z^h|_0 \leq Ch^{-2} |z^h|_1^2 \leq Ch^{-4} |z^h|_0^2 \quad \forall z^h \in S^h. \end{aligned} \quad (2.26)$$

LEMMA. 2.1 *Let the assumptions (A) hold. Then for all $z^h \in S^h$ we have that*

$$|z^h|_{1,s} \leq C |\Delta^h z^h|_0, \quad \text{for any } s \in (1, \infty), \quad (2.27a)$$

$$|z^h|_{1,4} \leq C |\Delta^h z^h|_0^{\frac{1}{2}} |z^h|_1^{\frac{1}{2}}. \quad (2.27b)$$

Furthermore

$$|\Delta^h(\pi^h \eta)|_0 \leq C |\eta|_2 \quad \forall \eta \in H^2(\Omega) \text{ with } \frac{\partial \eta}{\partial \nu} = 0 \text{ on } \partial\Omega. \quad (2.28)$$

Proof. The proof of (2.27a) can be found in Barrett, Langdon, and Nürnberg (2004, Lemma 3.1). However, we state the proof for the reader’s convenience as the proof of (2.27b) is very similar. It follows from (2.25) and (2.23) that

$$(I - \mathcal{f})z^h = -\mathcal{G}^h[\Delta^h z^h] \quad \forall z^h \in S^h. \quad (2.29)$$

For $s \in (2, \infty)$ we have from (2.29), (1.18), (2.15), (2.14), (2.22) and (2.24) that

$$\begin{aligned} |z^h|_{1,s} &\leq |\mathcal{G}[\Delta^h z^h]|_{1,s} + |(I - \pi^h)\mathcal{G}[\Delta^h z^h]|_{1,s} + |(\pi^h \mathcal{G} - \mathcal{G}^h)\Delta^h z^h|_{1,s} \\ &\leq C \|\mathcal{G}[\Delta^h z^h]\|_2 + Ch^{-(1-\frac{2}{s})} |(\pi^h \mathcal{G} - \mathcal{G}^h)\Delta^h z^h|_1 \leq C |\Delta^h z^h|_0 \quad \forall z^h \in S^h. \end{aligned}$$

Hence the desired result (2.27a), for all stated s , follows immediately. Similarly, we have from (2.29), (1.18), (2.15), (2.14), (2.22) and (2.24) that

$$\begin{aligned} |z^h|_{1,4} &\leq |\mathcal{G}[\Delta^h z^h]|_{1,4} + |(I - \pi^h)\mathcal{G}[\Delta^h z^h]|_{1,4} + |(\pi^h \mathcal{G} - \mathcal{G}^h)\Delta^h z^h|_{1,4} \\ &\leq |\mathcal{G}[\Delta^h z^h]|_1^{\frac{1}{2}} \|\mathcal{G}[\Delta^h z^h]\|_2^{\frac{1}{2}} + C h^{\frac{1}{2}} |\mathcal{G}[\Delta^h z^h]|_2 + C h^{-\frac{1}{2}} |(\pi^h \mathcal{G} - \mathcal{G}^h)\Delta^h z^h|_1 \\ &\leq |\mathcal{G}[\Delta^h z^h]|_1^{\frac{1}{2}} |\Delta^h z^h|_0^{\frac{1}{2}} + C h^{\frac{1}{2}} |\Delta^h z^h|_0 \quad \forall z^h \in S^h. \end{aligned} \quad (2.30)$$

It follows from (2.29), (2.24) and (2.26) that for all $z^h \in S^h$

$$|\mathcal{G}[\Delta^h z^h]|_1 \leq |\mathcal{G}^h[\Delta^h z^h]|_1 + |(\mathcal{G} - \mathcal{G}^h)[\Delta^h z^h]|_1 \leq |z^h|_1 + C h |\Delta^h z^h|_0 \leq C |z^h|_1.$$

Combining (2.30) and (2.26) yields that

$$|z^h|_{1,4} \leq C |z^h|_1^{\frac{1}{2}} |\Delta^h z^h|_0^{\frac{1}{2}} + C h^{\frac{1}{2}} |\Delta^h z^h|_0 \leq C |z^h|_1^{\frac{1}{2}} |\Delta^h z^h|_0^{\frac{1}{2}} \quad \forall z^h \in S^h;$$

and hence the desired result (2.27b).

Finally, it follows from (2.3), (2.17), (2.25), (2.15), $\eta \in H^2(\Omega)$ with $\frac{\partial \eta}{\partial \nu} = 0$ on $\partial\Omega$, and (2.13) that

$$\begin{aligned} |\Delta^h(\pi^h \eta)|_0^2 &\leq |\Delta^h(\pi^h \eta)|_h^2 = -(\nabla(\pi^h \eta), \nabla(\Delta^h(\pi^h \eta))) \\ &= -(\nabla \eta, \nabla(\Delta^h(\pi^h \eta))) + (\nabla(I - \pi^h)\eta, \nabla(\Delta^h(\pi^h \eta))) \\ &\leq |\Delta \eta|_0 |\Delta^h(\pi^h \eta)|_0 + C h |\eta|_2 |\nabla(\Delta^h(\pi^h \eta))|_0 \leq C |\eta|_2^2; \end{aligned}$$

and hence the desired result (2.28). \square

Similarly to (2.25), we introduce $L^h : \underline{S}^h \rightarrow \widehat{S}^h$ such that

$$(L^h \underline{z}^h, \underline{\chi}) = -(\mathcal{C} \underline{\mathcal{E}}(\underline{z}^h), \underline{\mathcal{E}}(\underline{\chi})) \quad \forall \underline{\chi} \in \underline{S}^h. \quad (2.31)$$

We introduce also $N_c : \underline{X} \rightarrow \widehat{H}^1(\Omega)$ and $N_c^h : \underline{X} \rightarrow \widehat{S}^h$, where $\underline{X} := \{\eta \in (\underline{H}^1(\Omega))' : \langle \eta, \underline{v} \rangle = 0 \quad \forall \underline{v} \in \underline{RM}\}$, such that

$$(\mathcal{C} \underline{\mathcal{E}}(N_c \underline{\xi}), \underline{\mathcal{E}}(\underline{\eta})) = \langle \underline{\xi}, \underline{\eta} \rangle \quad \forall \underline{\eta} \in \underline{H}^1(\Omega), \quad (2.32)$$

$$(\mathcal{C} \underline{\mathcal{E}}(N_c^h \underline{\xi}), \underline{\mathcal{E}}(\underline{\chi})) = \langle \underline{\xi}, \underline{\chi} \rangle \quad \forall \underline{\chi} \in \underline{S}^h. \quad (2.33)$$

As \mathcal{C} satisfies (1.9) and (1.10), the well-posedness of these operators is easily demonstrated. As Ω is convex polygonal, we will assume the analogue of (2.22),

$$\|N_c \underline{\xi}\|_2 \leq C |\underline{\xi}|_0 \quad \forall \underline{\xi} \in \underline{L}^2(\Omega) \cap \underline{X}. \quad (2.34)$$

If \mathcal{C} is isotropic, (1.11), then the singularity exponents in $N_c \underline{\xi}$ do not depend on the Lamé moduli; and (2.34) follows immediately, for example, on combining Grisvard (1989, Theorem I) and Seif (1973, Lemma 3.2). Unfortunately, if \mathcal{C} is anisotropic then the singularity exponents depend on the specific form of \mathcal{C} and there is no general result of the type (2.34) in the literature. However, there is also no counterexample. For any particular material law, \mathcal{C} , and domain Ω the singularity exponents in $N_c \underline{\xi}$ can be computed, see e.g. Costabel, Dauge, and Lafranche (2001), and hence the assumption (2.34) can be tested.

We now have the analogues of (2.26), (2.27a) and (2.28).

LEMMA. 2.2 *Let the assumptions (A) hold and, if \mathcal{C} is anisotropic, assume that (2.34) holds. Then for all $s \in (1, \infty)$ and for all $\underline{z}^h \in \underline{S}^h$ we have that*

$$|\underline{\underline{\mathcal{E}}}(\underline{z}^h)|_{0,s} \leq C |L^h \underline{z}^h|_0 \leq C h^{-1} |\underline{z}^h|_1. \quad (2.35)$$

Furthermore

$$|L^h(\pi^h \underline{\eta})|_0 \leq C |\underline{\eta}|_2 \quad \forall \underline{\eta} \in \underline{H}^2(\Omega) \quad \text{with} \quad \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{\eta}) \underline{\nu} = \underline{0} \text{ on } \partial\Omega. \quad (2.36)$$

Proof. It follows from (1.21), (1.10), (2.32), (2.33), (2.15) and (2.34) that

$$\begin{aligned} C \|(N_c - N_c^h) \underline{\xi}\|_1^2 &\leq C |\underline{\underline{\mathcal{E}}}((N_c - N_c^h) \underline{\xi})|_0^2 \leq (C \underline{\underline{\mathcal{E}}}((N_c - N_c^h) \underline{\xi}), \underline{\underline{\mathcal{E}}}((N_c - N_c^h) \underline{\xi})) \\ &= (C \underline{\underline{\mathcal{E}}}((N_c - N_c^h) \underline{\xi}), \underline{\underline{\mathcal{E}}}((I - \pi^h)(N_c \underline{\xi}))) \leq C |\underline{\underline{\mathcal{E}}}((I - \pi^h)(N_c \underline{\xi}))|_0^2 \\ &\leq C |(I - \pi^h)(N_c \underline{\xi})|_1^2 \leq C h^2 \|N_c \underline{\xi}\|_2^2 \leq C h^2 |\underline{\xi}|_0^2 \quad \forall \underline{\xi} \in \underline{L}^2(\Omega) \cap \underline{X}. \end{aligned} \quad (2.37)$$

Let $\underline{z}^h = (\underline{z}^h - \underline{z}_{\text{RM}}^h) + \underline{z}_{\text{RM}}^h$ such that $\underline{z}_{\text{RM}}^h \in \underline{\text{RM}}$ and $\underline{z}^h - \underline{z}_{\text{RM}}^h \in \widehat{S}^h$. Then it follows from (2.33) and (1.21) that

$$\underline{z}^h - \underline{z}_{\text{RM}}^h = -N_c^h (L^h \underline{z}^h). \quad (2.38)$$

Combining (2.38), (1.18), (2.16), (2.14) and (2.37) yields for $s \in (2, \infty)$

$$\begin{aligned} |\underline{\underline{\mathcal{E}}}(\underline{z}^h)|_{0,s} &= |\underline{\underline{\mathcal{E}}}(N_c^h (L^h \underline{z}^h))|_{0,s} \leq |\underline{\underline{\mathcal{E}}}(N_c (L^h \underline{z}^h))|_{0,s} + |\underline{\underline{\mathcal{E}}}((I - \pi^h)N_c (L^h \underline{z}^h))|_{0,s} \\ &\quad + |\underline{\underline{\mathcal{E}}}(\pi^h [N_c (L^h \underline{z}^h)] - N_c^h (L^h \underline{z}^h))|_{0,s} \\ &\leq C \|N_c (L^h \underline{z}^h)\|_2 + C h^{-(1-\frac{2}{s})} |\pi^h [N_c (L^h \underline{z}^h)] - N_c^h (L^h \underline{z}^h)|_1 \leq C |L^h \underline{z}^h|_0 \end{aligned}$$

and hence the first inequality in (2.35).

It follows from (2.31) and (2.13) that

$$\begin{aligned} |L^h \underline{z}^h|_0^2 &= -(C \underline{\underline{\mathcal{E}}}(\underline{z}^h), \underline{\underline{\mathcal{E}}}(L^h \underline{z}^h)) \leq C |\underline{\underline{\mathcal{E}}}(\underline{z}^h)|_0 |\underline{\underline{\mathcal{E}}}(L^h \underline{z}^h)|_0 \\ &\leq C h^{-1} |\underline{\underline{\mathcal{E}}}(\underline{z}^h)|_0 |L^h \underline{z}^h|_0 \leq C h^{-2} |\underline{\underline{\mathcal{E}}}(\underline{z}^h)|_0^2 \leq C h^{-2} |\underline{z}^h|_1^2 \end{aligned}$$

and hence the second inequality in (2.35).

Finally, it follows from (2.3), (2.17), (2.31), (2.15), $\underline{\eta} \in \underline{H}^2(\Omega)$ with $\mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{\eta}) \underline{\nu} = \underline{0}$ on $\partial\Omega$, and (2.13) that

$$\begin{aligned} |L^h(\pi^h \underline{\eta})|_0^2 &\leq |L^h(\pi^h \underline{\eta})|_h^2 = -(C \underline{\underline{\mathcal{E}}}(\pi^h \underline{\eta}), \underline{\underline{\mathcal{E}}}(L^h(\pi^h \underline{\eta}))) \\ &= -(C \underline{\underline{\mathcal{E}}}(\underline{\eta}), \underline{\underline{\mathcal{E}}}(L^h(\pi^h \underline{\eta}))) + (C \underline{\underline{\mathcal{E}}}((I - \pi^h)\underline{\eta}), \underline{\underline{\mathcal{E}}}(L^h(\pi^h \underline{\eta}))) \\ &\leq C |\underline{\eta}|_2 |L^h(\pi^h \underline{\eta})|_0 + C h |\underline{\eta}|_2 |\nabla(L^h(\pi^h \underline{\eta}))|_0 \leq C |\underline{\eta}|_2^2. \end{aligned}$$

Hence the desired result (2.36). \square

We introduce the projection operator $P^h : \underline{W}^{1,1}(\Omega) \rightarrow \widehat{S}^h$ such that

$$(\underline{\underline{\mathcal{E}}}(z - P^h z), \underline{\underline{\mathcal{E}}}(\chi)) = 0 \quad \forall \chi \in \widehat{S}^h. \quad (2.39)$$

It is crucial for our analysis to prove the following result.

LEMMA. 2.3 *Let the assumptions (A) hold and let $\delta \in \mathbb{R}_{>0}$ be as defined in Lemma 1.1. Then there exists $h_0 \in \mathbb{R}_{>0}$ and a $\widehat{\beta} \in C([\frac{2+\delta}{1+\delta}, \infty))$ such that for all $p \in [\frac{2+\delta}{1+\delta}, \infty)$ and for all $h \in (0, h_0)$*

$$|\underline{\mathcal{E}}(P^h \underline{z})|_{0,p} \leq \widehat{\beta}(p) |\underline{\mathcal{E}}(\underline{z})|_{0,p} \quad \forall \underline{z} \in \widehat{V}_p \quad (2.40)$$

with $\widehat{\beta}(p) \geq 1$ and $\widehat{\beta}(p) \rightarrow \widehat{\beta}(2) = 1$ as $p \rightarrow 2$.

Proof. We adapt the argument for the Laplacian with homogeneous Dirichlet boundary conditions given in Brenner and Scott (2002, Chapter 8), which is based on the approach in Rannacher and Scott (1982). Given \mathcal{T}^h and any $y \in \Omega$, let $\sigma_y \in \mathcal{T}^h$ be such that $y \in \sigma_y$. We then introduce $\delta_y^h \in C_0^\infty(\overline{\Omega})$ with $\text{supp}(\delta_y^h) \subset \sigma_y$ such that

$$\int_{\sigma_y} \delta_y^h dx = 1 \quad \text{and} \quad \|\delta_y^h\|_{m,\infty,\sigma_y} \leq C h^{-(2+m)} \quad \forall m \in \mathbb{N}. \quad (2.41)$$

For $i, j \in \{1, 2\}$, let $\underline{f}_{y,ij} \in \widehat{H}^1(\Omega)$ be such that

$$(\underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(\underline{\eta})) = (\delta_y^h, [\underline{\mathcal{E}}(\underline{\eta})]_{ij}) \quad \forall \underline{\eta} \in \underline{H}^1(\Omega). \quad (2.42)$$

It follows from (1.14) and (1.21) that (2.42) is well-posed. We have from (2.41), (2.39) and (2.42) for all $y \in \Omega$ and for $i, j \in \{1, 2\}$ that

$$\begin{aligned} |\underline{\mathcal{E}}(P^h \underline{z})]_{ij}(y) &= (\delta_y^h, [\underline{\mathcal{E}}(P^h \underline{z})]_{ij}) = (\underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(P^h \underline{z})) = (\underline{\mathcal{E}}(P^h \underline{f}_{y,ij}), \underline{\mathcal{E}}(\underline{z})) \\ &= (\delta_y^h, [\underline{\mathcal{E}}(\underline{z})]_{ij}) + (\underline{\mathcal{E}}([P^h - I]\underline{f}_{y,ij}), \underline{\mathcal{E}}(\underline{z})) \quad \forall \underline{z} \in \underline{H}^1(\Omega). \end{aligned} \quad (2.43)$$

For any $y \in \Omega$ and any constant $\rho \geq 1$, we introduce the weight function

$$\omega_{y,\rho}(x) := (|x - y|^2 + \rho^2 h^2)^{\frac{1}{2}}. \quad (2.44)$$

It is easily verified for any $\alpha \in \mathbb{R}$ that

$$\max_{\sigma \in \mathcal{T}^h} (\sup_{x \in \sigma} [\omega_{y,\rho}(x)]^\alpha / \inf_{x \in \sigma} [\omega_{y,\rho}(x)]^\alpha) \leq C, \quad |\omega_{y,\rho}^\alpha|_{0,\infty} \leq C \max\{1, (\rho h)^\alpha\} \quad (2.45a)$$

$$\text{and} \quad \left| \frac{\partial^m}{\partial x_i^m} [\omega_{y,\rho}(x)]^\alpha \right| \leq C(\alpha) [\omega_{y,\rho}(x)]^{\alpha-m} \quad \forall x \in \Omega, \quad \forall m \in \mathbb{N}, \quad i \in \{1, 2\}; \quad (2.45b)$$

where the positive constant $C(\alpha)$ depends continuously on α and is independent of the choice of $y \in \Omega$ and $\rho \geq 1$. It follows immediately from (2.15) and (2.45a) that for all $\sigma \in \mathcal{T}^h$, $\alpha \in \mathbb{R}$, $m \in \{0, 1\}$ and $i \in \{1, 2\}$

$$\int_{\sigma} \omega_{y,\rho}^\alpha \left(\frac{\partial^m}{\partial x_i^m} [(I - \pi^h)\eta] \right)^2 dx \leq C h^{2(2-m)} \int_{\sigma} \omega_{y,\rho}^\alpha \left[\left(\frac{\partial^2 \eta}{\partial x_1^2} \right)^2 + \left(\frac{\partial^2 \eta}{\partial x_1 \partial x_2} \right)^2 + \left(\frac{\partial^2 \eta}{\partial x_2^2} \right)^2 \right] dx \quad \forall \eta \in H^2(\sigma). \quad (2.46)$$

It follows from (2.43), a Hölder inequality and (2.45a) that for any $p \in (2, \infty)$, $\alpha > 0$ and $\rho \geq 1$

$$\begin{aligned} |\underline{\underline{\mathcal{E}}}(P^h \underline{z})|_{0,p} &\leq C \left[1 + \left(\sup_{y \in \Omega} \int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)} dx \right)^{\frac{1}{2}} M_{\rho,\alpha}^h \right] |\underline{\underline{\mathcal{E}}}(z)|_{0,p} \\ &\leq C \left[1 + \alpha^{-\frac{1}{2}} (\rho h)^{-\frac{\alpha}{2}} M_{\rho,\alpha}^h \right] |\underline{\underline{\mathcal{E}}}(z)|_{0,p} \quad \forall z \in \widehat{V}_p, \end{aligned} \quad (2.47)$$

$$\text{where} \quad M_{\rho,\alpha}^h := \max_{i,j=1,2} \sup_{y \in \Omega} \left\{ \int_{\Omega} \omega_{y,\rho}^{\alpha+2} |\underline{\underline{\mathcal{E}}}([I - P^h] f_{y,ij})|^2 dx \right\}^{\frac{1}{2}}. \quad (2.48)$$

The goal is to prove the analogue of Brenner and Scott (2002, Lemma 8.2.6); that is, for appropriate $\alpha > 0$ and ρ sufficiently large that there exists an h_0 such that

$$M_{\rho,\alpha}^h \leq C h^{\frac{\alpha}{2}} \quad \forall h \in (0, h_0). \quad (2.49)$$

It would then follow from (2.47) and (2.49) that (2.40) holds with $\widehat{\beta}(p) = C_1$ for all $p \in (2, \infty)$, for some constant C_1 . In addition, it would follow from (1.22), (2.39) and the above bound for $p \in (2, \infty)$ that for $p \in [\frac{2+\delta}{1+\delta}, 2)$ and for all $z \in \widehat{V}_p$

$$|\underline{\underline{\mathcal{E}}}(P^h \underline{z})|_{0,p} \leq \beta(p) \sup_{\underline{0} \neq \underline{\eta} \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(P^h \underline{z}), \underline{\underline{\mathcal{E}}}(\underline{\eta}))}{|\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,q}} = \beta(p) \sup_{\underline{0} \neq \underline{\eta} \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(z), \underline{\underline{\mathcal{E}}}(P^h \underline{\eta}))}{|\underline{\underline{\mathcal{E}}}(\underline{\eta})|_{0,q}} \leq \beta(p) C_1 |\underline{\underline{\mathcal{E}}}(z)|_{0,p}, \quad (2.50)$$

where $\frac{1}{p} + \frac{1}{q} = 1$. As (2.40) trivially holds with $\widehat{\beta}(2) = 1$ from inspecting (2.39), it follows that (2.40) holds with $\widehat{\beta}(p) = C_2$ for all $p \in [\frac{2+\delta}{1+\delta}, \infty)$, for some constant C_2 . Moreover, the desired result (2.40) holds for all $p \in [\frac{2+\delta}{1+\delta}, \infty)$ by applying the Riesz–Thorin theorem as in Lemma 1.1 to the P^h induced mapping that takes $\underline{\underline{\mathcal{E}}}(z) \in [L^p(\Omega)]_{\text{sym}}^{2 \times 2}$ to $\underline{\underline{\mathcal{E}}}(P^h \underline{z}) \in [L^p(\Omega)]_{\text{sym}}^{2 \times 2}$.

Therefore we need to prove (2.49). Let $\underline{\mathcal{Y}} := \{ \{ \underline{\xi}, \underline{\zeta} \} \in (\underline{H}^1(\Omega))' \times \underline{L}^2(\partial\Omega) : \langle \underline{\xi}, \underline{v} \rangle + \int_{\partial\Omega} \underline{\zeta} \cdot \underline{v} ds = 0 \quad \forall \underline{v} \in \underline{\mathbf{RM}} \}$. Then $N : \underline{\mathcal{Y}} \rightarrow \widehat{\underline{H}}^1(\Omega)$ is such that

$$(\underline{\underline{\mathcal{E}}}(N(\underline{\xi}, \underline{\zeta})), \underline{\underline{\mathcal{E}}}(\underline{\eta})) = \langle \underline{\xi}, \underline{\eta} \rangle + \int_{\partial\Omega} \underline{\zeta} \cdot \underline{\eta} ds \quad \forall \underline{\eta} \in \underline{H}^1(\Omega). \quad (2.51)$$

Let $\partial\Omega \equiv \cup_{j=1}^{J_B} \overline{\partial_j \Omega}$ and $\partial_j \Omega \cap \partial_k \Omega = \emptyset$ for $j \neq k$; with $\underline{\nu}^{(j)}$ the outward unit normal to $\partial_j \Omega$. In addition, let the largest inner angle ω of the convex polygonal domain Ω be such that $\omega \leq \frac{r}{2r-1} \pi$ for some $r > 1$. Then, similarly to (2.34), on combining Grisvard (1989, Theorem I), $\varphi(z) := \sin^2(\omega z) - z^2 \sin \omega \Rightarrow \varphi(iz) = \widetilde{\varphi}(z) := z^2 \sin^2 \omega - \sinh^2(\omega z)$, and the fact that $\widetilde{\varphi}(z)$ has no roots such that $|\text{Im}(z)| \leq \frac{\pi}{\omega}$, apart from the double root at $z = 0$ and the simple roots at $z = \pm i$, see Seif (1973, Lemma 3.2); we have for $p \in (1, 2r]$ that

$$\|N(\underline{\xi}, \underline{\zeta})\|_{2,p} \leq C \left[\|\underline{\xi}\|_{0,p} + \sum_{j=1}^{J_B} \|\underline{\zeta}\|_{1-\frac{1}{p}, p, \partial_j \Omega} \right] \quad \forall \{ \underline{\xi}, \underline{\zeta} \} \in (\underline{L}^p(\Omega) \times \prod_{j=1}^{J_B} \underline{W}^{1-\frac{1}{p}, p}(\partial_j \Omega)) \cap \underline{\mathcal{Y}}; \quad (2.52)$$

provided that the compatibility condition, Grisvard (1989, (1.5)),

$$\underline{\zeta} |_{\partial_j \Omega} \cdot \underline{\nu}^{(j+1)} = \underline{\zeta} |_{\partial_{j+1} \Omega} \cdot \underline{\nu}^{(j)} \quad \text{at every vertex } S_j \text{ of } \Omega \quad (2.53)$$

holds (in the integral sense if $p = 2$).

For fixed $y \in \Omega$ and $i, j \in \{1, 2\}$, let $\underline{e} := (I - P^h) \underline{f}_{y,ij} \in \widehat{H}^1(\Omega)$, $\underline{e}^A := (I - \pi^h) \underline{f}_{y,ij} \in \underline{H}^1(\Omega)$ and $\underline{e}^h := (\pi^h - P^h) \underline{f}_{y,ij} \in \underline{S}^h$. We note that

$$\underline{\mathcal{E}}(\eta \underline{z}) = \eta \underline{\mathcal{E}}(\underline{z}) + \frac{1}{2} [\underline{z} \otimes (\nabla \eta) + (\nabla \eta) \otimes \underline{z}], \quad (2.54)$$

where $\underline{a} \otimes \underline{b} := \underline{a} \underline{b}^T$ for all $\underline{a}, \underline{b} \in \mathbb{R}^2$. It then follows from (2.54), (2.39), (1.25), (2.45a,b), (2.46) and (2.13) for any $y \in \Omega$, $i, j \in \{1, 2\}$, $\alpha > 0$ and $\rho \geq 1$ that

$$\begin{aligned} (\omega_{y,\rho}^{\alpha+2} \underline{\mathcal{E}}(\underline{e}), \underline{\mathcal{E}}(\underline{e})) &= (\underline{\mathcal{E}}(\underline{e}), \underline{\mathcal{E}}(\omega_{y,\rho}^{\alpha+2} \underline{e}^A) + \underline{\mathcal{E}}((I - \pi^h)[\omega_{y,\rho}^{\alpha+2} \underline{e}^h])) \\ &\quad - \frac{1}{2} (\underline{\mathcal{E}}(\underline{e}), [\underline{e} \otimes (\nabla \omega_{y,\rho}^{\alpha+2}) + (\nabla \omega_{y,\rho}^{\alpha+2}) \otimes \underline{e}]) \\ &\leq C(\alpha) \left[\int_{\Omega} [\omega_{y,\rho}^{\alpha+2} |\underline{\mathcal{E}}(\underline{e}^A)|^2 + \omega_{y,\rho}^{\alpha} |\underline{e}^A|^2] dx + \int_{\Omega} \omega_{y,\rho}^{\alpha} |\underline{e}^h|^2 dx \right. \\ &\quad \left. + \int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)} |\underline{\mathcal{E}}((I - \pi^h)[\omega_{y,\rho}^{\alpha+2} \underline{e}^h])|^2 dx \right] \\ &\leq C(\alpha) \left[\int_{\Omega} [\omega_{y,\rho}^{\alpha+2} |\underline{\mathcal{E}}(\underline{e}^A)|^2 + \omega_{y,\rho}^{\alpha} |\underline{e}^A|^2] dx + \int_{\Omega} \omega_{y,\rho}^{\alpha} |\underline{e}^h|^2 dx \right]. \end{aligned} \quad (2.55)$$

Let $\underline{\psi} = N((I - P_{\text{RM}})(\omega_{y,\rho}^{\alpha} \underline{e}), \underline{0})$, where $P_{\text{RM}} : \underline{L}^2(\Omega) \rightarrow \underline{\text{RM}}$ is such that

$$((I - P_{\text{RM}})\underline{z}, \underline{\eta}) = 0 \quad \forall \underline{\eta} \in \underline{\text{RM}}. \quad (2.56)$$

It follows from (1.21), (1.14) and (2.56) that

$$\begin{aligned} |\underline{v}|_1 &\leq |(I - P_{\text{RM}})\underline{v}|_1 + |P_{\text{RM}}\underline{v}|_1 \leq C |\underline{\mathcal{E}}((I - P_{\text{RM}})\underline{v})|_0 + |P_{\text{RM}}\underline{v}|_1 \\ &\leq C [|\underline{\mathcal{E}}(\underline{v})|_0 + |P_{\text{RM}}\underline{v}|_0] \leq C [|\underline{\mathcal{E}}(\underline{v})|_0 + |\underline{v}|_0] \quad \forall \underline{v} \in \underline{H}^1(\Omega). \end{aligned} \quad (2.57)$$

We have, on noting (2.51) and (2.39), that for all $\varsigma > 0$

$$\begin{aligned} (\omega_{y,\rho}^{\alpha} \underline{e}, \underline{e}) &= (\underline{\mathcal{E}}(\underline{\psi}), \underline{\mathcal{E}}(\underline{e})) = (\underline{\mathcal{E}}((I - \pi^h)\underline{\psi}), \underline{\mathcal{E}}(\underline{e})) \\ &\leq \varsigma (\omega_{y,\rho}^{\alpha+2} \underline{\mathcal{E}}(\underline{e}), \underline{\mathcal{E}}(\underline{e})) + C \varsigma^{-1} \int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)} |\underline{\mathcal{E}}((I - \pi^h)\underline{\psi})|^2 dx. \end{aligned} \quad (2.58)$$

It follows from (2.46) and (2.44) that

$$\begin{aligned} \int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)} |\underline{\mathcal{E}}((I - \pi^h)\underline{\psi})|^2 dx &\leq C h^2 \sum_{k,\ell=1}^2 \int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)} \left| \frac{\partial^2 \underline{\psi}}{\partial x_k \partial x_\ell} \right|^2 dx \\ &\leq C h^2 \left(\int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)r'} dx \right)^{\frac{1}{r'}} \|\underline{\psi}\|_{2,2r}^2 \leq C(\alpha) \rho^{-2} (\rho h)^{\frac{2}{r'} - \alpha} \|\underline{\psi}\|_{2,2r}^2, \end{aligned} \quad (2.59)$$

where r is as defined in (2.52) and $\frac{1}{r} + \frac{1}{r'} = 1$. Next we note that (2.52), (1.18), (1.21), (2.44), (2.54) and (2.45b) yield, on assuming that $\alpha \in (0, \frac{2(r-1)}{r})$,

$$\begin{aligned} \|\underline{\psi}\|_{2,2r}^2 &\leq C \|(I - P_{\text{RM}})(\omega_{y,\rho}^\alpha \underline{e})\|_{0,2r}^2 \leq C \|(I - P_{\text{RM}})(\omega_{y,\rho}^\alpha \underline{e})\|_{1,\frac{2r}{r+1}}^2 \\ &\leq C \|\underline{\mathcal{E}}(\omega_{y,\rho}^\alpha \underline{e})\|_{0,\frac{2r}{r+1}}^2 \leq C \left(\int_{\Omega} \omega_{y,\rho}^{(\alpha-2)r} dx \right)^{\frac{1}{r}} \int_{\Omega} \omega_{y,\rho}^{2-\alpha} |\underline{\mathcal{E}}(\omega_{y,\rho}^\alpha \underline{e})|^2 dx \\ &\leq C(\alpha) (\rho h)^{\alpha-2+\frac{2}{r}} [(\omega_{y,\rho}^{\alpha+2} \underline{\mathcal{E}}(\underline{e}), \underline{\mathcal{E}}(\underline{e})) + (\omega_{y,\rho}^\alpha \underline{e}, \underline{e})]. \end{aligned} \quad (2.60)$$

Therefore for any fixed $\alpha \in (0, \frac{2(r-1)}{r})$, we have for all $y \in \Omega$, $i, j \in \{1, 2\}$, $\rho > \rho_0(\alpha)$ and $h > 0$ on combining (2.55), (2.58) with ς sufficiently small, (2.59) and (2.60) that

$$(\omega_{y,\rho}^{\alpha+2} \underline{\mathcal{E}}(\underline{e}), \underline{\mathcal{E}}(\underline{e})) \leq C(\alpha, \rho) \int_{\Omega} [\omega_{y,\rho}^{\alpha+2} |\underline{\mathcal{E}}(\underline{e}^A)|^2 + \omega_{y,\rho}^\alpha |\underline{e}^A|^2] dx. \quad (2.61)$$

Hence the desired result (2.49) follows from (2.48), (2.61) and (2.46); if we can show for any $y \in \Omega$, $i, j \in \{1, 2\}$, $\alpha \in (0, 1)$, $\rho \geq 1$ and $h > 0$ that

$$\max_{k,\ell=1,2} \int_{\Omega} \omega_{y,\rho}^{\alpha+2} \left| \frac{\partial^2 \underline{f}_{y,ij}}{\partial x_k \partial x_\ell} \right|^2 dx \leq C(\alpha, \rho) h^{\alpha-2}. \quad (2.62)$$

First, we have from (2.45b) that

$$\max_{k,\ell=1,2} \int_{\Omega} \omega_{y,\rho}^{\alpha+2} \left| \frac{\partial^2 \underline{f}_{y,ij}}{\partial x_k \partial x_\ell} \right|^2 dx \leq C(\alpha) \left[|\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{f}_{y,ij}|_2^2 + \int_{\Omega} [\omega_{y,\rho}^\alpha |\nabla \underline{f}_{y,ij}|^2 + \omega_{y,\rho}^{\alpha-2} |\underline{f}_{y,ij}|^2] dx \right]. \quad (2.63)$$

Second, it follows from (2.56), (1.14), (2.54), the symmetry of $\underline{\mathcal{E}}(\cdot)$ and (2.42) that $(I - P_{\text{RM}})(\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{f}_{y,ij}) \in \widehat{H}^1(\Omega)$ solves for all $\underline{\eta} \in \underline{H}^1(\Omega)$

$$\begin{aligned} (\underline{\mathcal{E}}((I - P_{\text{RM}})(\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{f}_{y,ij})), \underline{\mathcal{E}}(\underline{\eta})) &= (\underline{\mathcal{E}}(\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{f}_{y,ij}), \underline{\mathcal{E}}(\underline{\eta})) \\ &= (\underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{\eta})) + \frac{1}{2} ([\underline{f}_{y,ij} \otimes \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) + \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) \otimes \underline{f}_{y,ij}], \underline{\mathcal{E}}(\underline{\eta})) \\ &\quad - (\underline{\mathcal{E}}(\underline{f}_{y,ij}) \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}), \underline{\eta}) \\ &= -\frac{1}{2} ([\underline{e}_i \otimes \underline{e}_j + \underline{e}_j \otimes \underline{e}_i] \nabla \delta_y^h, \omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{\eta}) \\ &\quad - (\underline{\mathcal{E}}(\underline{f}_{y,ij}) \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) + \frac{1}{2} \nabla \cdot [\underline{f}_{y,ij} \otimes \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) + \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) \otimes \underline{f}_{y,ij}], \underline{\eta}) \\ &\quad + \frac{1}{2} \int_{\partial\Omega} [[\underline{f}_{y,ij} \otimes \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) + \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) \otimes \underline{f}_{y,ij}] \underline{\nu}] \cdot \underline{\eta} ds. \end{aligned} \quad (2.64)$$

Noting (2.51) and (2.53), and applying the bounds (2.52), (2.45b) and the trace inequality $\|\cdot\|_{\frac{1}{2},\partial_k\Omega} \leq C \|\cdot\|_{1,\Omega}$ to (2.64) yields that

$$\begin{aligned} |\omega_{y,\rho}^{\frac{\alpha}{2}+1} \underline{f}_{y,ij}|_2 &\leq C [|\omega_{y,\rho}^{\frac{\alpha}{2}+1} \nabla \delta_y^h|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}} \nabla \underline{f}_{y,ij}|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0 \\ &\quad + \sum_{k=1}^{J_B} \|[\underline{f}_{y,ij} \otimes \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) + \nabla(\omega_{y,\rho}^{\frac{\alpha}{2}+1}) \otimes \underline{f}_{y,ij}] \underline{\nu}\|_{\frac{1}{2},\partial_k\Omega}] \\ &\leq C [|\omega_{y,\rho}^{\frac{\alpha}{2}+1} \nabla \delta_y^h|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}} \nabla \underline{f}_{y,ij}|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0]. \end{aligned} \quad (2.65)$$

It follows from (2.45b), (2.57), (2.45a) and (2.54) that

$$\begin{aligned} |\omega_{y,\rho}^{\frac{\alpha}{2}} \nabla \underline{f}_{y,ij}|_0 &\leq C \left[|\omega_{y,\rho}^{\frac{\alpha}{2}} \underline{f}_{y,ij}|_1 + |\nabla(\omega_{y,\rho}^{\frac{\alpha}{2}} \underline{f}_{y,ij})|_0 \right] \leq C \left[|\underline{\mathcal{E}}(\omega_{y,\rho}^{\frac{\alpha}{2}} \underline{f}_{y,ij})|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0 \right] \\ &\leq C \left[|\omega_{y,\rho}^{\frac{\alpha}{2}} \underline{\mathcal{E}}(\underline{f}_{y,ij})|_0 + |\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0 \right]. \end{aligned} \quad (2.66)$$

We have from (2.54) that

$$\begin{aligned} (\omega_{y,\rho}^{\alpha} \underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(\underline{f}_{y,ij})) &= (\underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(\omega_{y,\rho}^{\alpha} \underline{f}_{y,ij})) \\ &\quad - \frac{1}{2}(\underline{f}_{y,ij} \otimes \nabla(\omega_{y,\rho}^{\alpha}) + \nabla(\omega_{y,\rho}^{\alpha}) \otimes \underline{f}_{y,ij}, \underline{\mathcal{E}}(\underline{f}_{y,ij})). \end{aligned} \quad (2.67)$$

Similarly to (2.64), testing (2.42) with $\underline{\eta} = \omega_{y,\rho}^{\alpha} \underline{f}_{y,ij}$ yields that

$$(\underline{\mathcal{E}}(\underline{f}_{y,ij}), \underline{\mathcal{E}}(\omega_{y,\rho}^{\alpha} \underline{f}_{y,ij})) = -\frac{1}{2}([\underline{e}_i \otimes \underline{e}_j + \underline{e}_j \otimes \underline{e}_i] \nabla \delta_y^h, \omega_{y,\rho}^{\alpha} \underline{f}_{y,ij}). \quad (2.68)$$

Combining (2.63), (2.65), (2.66), (2.67) and (2.68) yields that

$$\max_{k,\ell=1,2} \int_{\Omega} \omega_{y,\rho}^{\alpha+2} \left| \frac{\partial^2 \underline{f}_{y,ij}}{\partial x_k \partial x_\ell} \right|^2 dx \leq C(\alpha) \left[|\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0^2 + |\omega_{y,\rho}^{\frac{\alpha}{2}+1} \nabla \delta_y^h|_0^2 \right]. \quad (2.69)$$

For $p \in (1, \frac{2}{\alpha})$, let $\underline{\Upsilon} = N((I - P_{\text{RM}})\underline{\xi}, \underline{0})$, where $[\underline{\xi}]_\ell = \text{sign}([\underline{f}_{y,ij}]_\ell) |[\underline{f}_{y,ij}]_\ell|^{2p-1}$, $\ell = 1, 2$. It follows from (2.56), (2.51), (2.42), (1.18), (2.52) and (2.44) that

$$\begin{aligned} |\underline{f}_{y,ij}|_{0,2p}^{2p} &= (\underline{\xi}, \underline{f}_{y,ij}) = ((I - P_{\text{RM}})\underline{\xi}, \underline{f}_{y,ij}) = (\underline{\mathcal{E}}(\underline{\Upsilon}), \underline{\mathcal{E}}(\underline{f}_{y,ij})) = (\delta_y^h, [\underline{\mathcal{E}}(\underline{\Upsilon})]_{ij}) \\ &\leq C |\delta_y^h|_{0, \frac{2p}{p+1}} |\underline{\Upsilon}|_{1, \frac{2p}{p-1}} \leq C |\delta_y^h|_{0, \frac{2p}{p+1}} \|\underline{\Upsilon}\|_{2, \frac{2p}{2p-1}} \leq C |\delta_y^h|_{0, \frac{2p}{p+1}} |(I - P_{\text{RM}})\underline{\xi}|_{0, \frac{2p}{2p-1}} \\ &\leq C |\delta_y^h|_{0, \frac{2p}{p+1}} |\underline{\xi}|_{0, \frac{2p}{2p-1}} \leq C |\delta_y^h|_{0, \frac{2p}{p+1}} |\underline{f}_{y,ij}|_{0,2p}^{2p-1} \leq C |\delta_y^h|_{0, \frac{2p}{p+1}}^{2p} \\ &\leq C \left(\int_{\Omega} \omega_{y,\rho}^{-(\alpha+2)p} dx \right) |\omega_{y,\rho}^{\frac{\alpha}{2}+1} \delta_y^h|_{0,2p}^{2p} \leq C(\alpha) (\rho h)^{2-(\alpha+2)p} |\omega_{y,\rho}^{\frac{\alpha}{2}+1} \delta_y^h|_{0,2p}^{2p}. \end{aligned} \quad (2.70)$$

Next we have from (2.44) and (2.70) that for $p \in (1, \frac{2}{\alpha})$

$$|\omega_{y,\rho}^{\frac{\alpha}{2}-1} \underline{f}_{y,ij}|_0^2 \leq C \left(\int_{\Omega} \omega_{y,\rho}^{(\alpha-2)p'} dx \right)^{\frac{1}{p'}} |\underline{f}_{y,ij}|_{0,2p}^2 \leq C(\alpha) (\rho h)^{\alpha-\frac{2}{p}} |\underline{f}_{y,ij}|_{0,2p}^2, \quad (2.71)$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. Finally, combining (2.69), (2.71), (2.70), (2.41) and (2.44) yields that

$$\begin{aligned} \max_{k,\ell=1,2} \int_{\Omega} \omega_{y,\rho}^{\alpha+2} \left| \frac{\partial^2 \underline{f}_{y,ij}}{\partial x_k \partial x_\ell} \right|^2 dx &\leq C(\alpha, \rho) h^{-2} |\omega_{y,\rho}^{\frac{\alpha}{2}+1} \delta_y^h|_0^2 \leq C(\alpha, \rho) h^{-6} |\omega_{y,\rho}^{\frac{\alpha}{2}+1}|_{0,\sigma_y}^2 \\ &\leq C(\alpha, \rho) h^{\alpha-2} \end{aligned}$$

and hence the desired result (2.62). \square

We now have a discrete analogue of a result similar to (1.22).

LEMMA. 2.4 *Let the assumptions of Lemma 2.3 hold. Then there exists $\delta_1 \in (0, \delta)$ and $C(c_0, m_c, M_c) \in \mathbb{R}_{>0}$ such that for all $p \in [2, 2 + \delta_1]$ and for all $h \in (0, h_0)$*

$$|\underline{\underline{\mathcal{E}}}(z^h)|_{0,p} \leq C \sup_{0 \neq \chi \in \widehat{S}^h} \frac{(c(\theta^h) \mathcal{C} \underline{\underline{\mathcal{E}}}(z^h), \underline{\underline{\mathcal{E}}}(\chi))}{|\underline{\underline{\mathcal{E}}}(\chi)|_{0,q}} \quad \forall z^h \in \widehat{S}^h, \quad \forall \theta^h \in K^h; \quad (2.72)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. The proof is an extension of the approach in Brenner and Scott (2002, §8.6) for a scalar second order linear elliptic equation. Similarly to (2.50), it follows from (1.22), (2.39) and (2.40) that for all $p \in [2, 2 + \delta]$, for all $h \in (0, h_0)$ and for all $z^h \in \widehat{S}^h$

$$|\underline{\underline{\mathcal{E}}}(z^h)|_{0,p} \leq \beta(p) \widehat{\beta}(q) \sup_{0 \neq \eta \in \widehat{V}_q} \frac{(\underline{\underline{\mathcal{E}}}(z^h), \underline{\underline{\mathcal{E}}}(P^h \eta))}{|\underline{\underline{\mathcal{E}}}(P^h \eta)|_{0,q}} \leq (1 + \sigma(p)) \sup_{0 \neq \chi \in \widehat{S}^h} \frac{(\underline{\underline{\mathcal{E}}}(z^h), \underline{\underline{\mathcal{E}}}(\chi))}{|\underline{\underline{\mathcal{E}}}(\chi)|_{0,q}}; \quad (2.73)$$

where $\sigma \in C([2, 2 + \delta])$, $\sigma(p) \geq 0$ and $\sigma(p) \rightarrow 0$ as $p \rightarrow 2$. On recalling (1.4) and (1.10) we define for all $z \in \widehat{V}_p$, for all $\eta \in \widehat{V}_q$ and for all $\theta^h \in K^h$

$$B(z, \eta) := (\mathcal{I} - \frac{1}{M_c} c(\theta^h) \mathcal{C}) \underline{\underline{\mathcal{E}}}(z), \underline{\underline{\mathcal{E}}}(\eta).$$

It follows from (1.10) and (1.4) that

$$|B(z, \eta)| \leq (1 - \frac{c_0 m_c}{M_c}) |\underline{\underline{\mathcal{E}}}(z)|_{0,p} |\underline{\underline{\mathcal{E}}}(\eta)|_{0,q}. \quad (2.74)$$

Combining (2.73) and (2.74) yields for all $z^h \in \widehat{S}^h$ and $\theta^h \in K^h$ that

$$\left[\frac{1}{1 + \sigma(p)} - (1 - \frac{c_0 m_c}{M_c}) \right] |\underline{\underline{\mathcal{E}}}(z^h)|_{0,p} \leq \frac{1}{M_c} \sup_{0 \neq \chi \in \widehat{S}^h} \frac{(c(\theta^h) \mathcal{C} \underline{\underline{\mathcal{E}}}(z^h), \underline{\underline{\mathcal{E}}}(\chi))}{|\underline{\underline{\mathcal{E}}}(\chi)|_{0,q}}. \quad (2.75)$$

Since $\sigma(p) \rightarrow 0$ as $p \rightarrow 2$ and σ is continuous, one can choose $\delta_1 \in (0, \delta)$ such that $\sigma(p) \leq \frac{1}{2} \frac{c_0 m_c}{M_c - c_0 m_c}$ for all $p \in [2, 2 + \delta_1]$. Hence (2.75) yields the desired result (2.72). \square

REMARK. 2.2 It is now straightforward to establish a global $L^\infty(0, T; \underline{W}^{1,p}(\Omega))$, $p > 2$, bound for \underline{u} solving (1.12e). Let $\theta \in L^\infty(\Omega_T)$. Then, similarly to the proof of Lemma 2.4, it follows from (1.21), (1.22), (1.10), (1.4), (1.12e) and a trace inequality that for *a.a.* $t \in (0, T)$

$$\left[\frac{1}{\beta(p)} - (1 - \frac{c_0 m_c}{M_c}) \right] |\underline{\underline{\mathcal{E}}}(\underline{u}(\cdot, t))|_{0,p} \leq \frac{1}{M_c} \sup_{0 \neq \eta \in \widehat{V}_q} \frac{|\int_{\partial\Omega} \underline{g} \cdot \underline{\eta} \, ds|}{|\underline{\underline{\mathcal{E}}}(\eta)|_{0,q}} \leq C \sup_{0 \neq \eta \in \widehat{V}_q} \frac{|\eta|_{0,1,\partial\Omega}}{\|\eta\|_{1,q}} \leq C. \quad (2.76)$$

We introduce for all $\varepsilon \in (0, 1)$, $b_\varepsilon : [-1, 1] \rightarrow [\varepsilon(2 - \varepsilon), 1]$ defined, on recalling (2.5), (2.7) and (2.8) by

$$b_\varepsilon(s) := \frac{1}{G''_\varepsilon(s)} \geq \frac{1}{G''(s)} = b(s). \quad (2.77)$$

Then the following two lemmas follow immediately from the construction of Ξ_ε , see Barrett, Nürnberg, and Styles (2004, Lemmas 2.2 and 2.3) for details.

LEMMA. 2.5 *Let the assumptions (A) hold. Then for any given $\varepsilon \in (0, 1)$ the function $\Xi_\varepsilon : S^h \rightarrow [L^\infty(\Omega)]^{2 \times 2}$ satisfies for all $z^h \in K^h$, $\underline{\xi} \in \mathbb{R}^2$ and for all $\sigma \in \mathcal{T}^h$*

$$\varepsilon (2 - \varepsilon) \underline{\xi}^T \underline{\xi} \leq \min_{x \in \bar{\sigma}} b_\varepsilon(z^h(x)) \underline{\xi}^T \underline{\xi} \leq \underline{\xi}^T \Xi_\varepsilon(z^h)|_\sigma \underline{\xi} \leq \max_{x \in \bar{\sigma}} b_\varepsilon(z^h(x)) \underline{\xi}^T \underline{\xi} \leq \underline{\xi}^T \underline{\xi}. \quad (2.78)$$

LEMMA. 2.6 *Let the assumptions (A) hold and let $\|\cdot\|$ denote the spectral norm on $\mathbb{R}^{2 \times 2}$. Then for any given $\varepsilon \in (0, 1)$ the function $\Xi_\varepsilon : S^h \rightarrow [L^\infty(\Omega)]^{2 \times 2}$ is such that for all $z^h \in K^h$ and for all $\sigma \in \mathcal{T}^h$*

$$\max_{x \in \bar{\sigma}} \|\Xi_\varepsilon(z^h) - b_\varepsilon(z^h) \mathcal{I}\|(x) \leq h_\sigma |\nabla[b_\varepsilon(z^h)]|_{0, \infty, \sigma} \leq 2 h_\sigma |\nabla z^h|_\sigma. \quad (2.79)$$

In the remainder of this section, we establish stability bounds for the solution of (2.12a–c) that are needed for our convergence analysis in §3.

LEMMA. 2.7 *Let the assumptions (A) hold and $\Theta_\varepsilon^{n-1} \in K^h$. Then for all $\varepsilon \in (0, 1)$ and for all $h, \tau_n > 0$ there exists a solution $\{\underline{U}_\varepsilon^n, \Theta_\varepsilon^n, W_\varepsilon^n\}$ to the n -th step of $(P_\varepsilon^{h, \tau})$ with $\mathcal{f}\Theta_\varepsilon^n = \mathcal{f}\Theta_\varepsilon^{n-1}$. $\{\underline{U}_\varepsilon^n, \Theta_\varepsilon^n\}$ is unique. In addition, W_ε^n is unique if there exists $j \in J$ such that $\Theta_\varepsilon^n(p_j) \in (-1, 1)$. Moreover, it holds that*

$$\begin{aligned} \mathcal{J}(\Theta_\varepsilon^n, \underline{U}_\varepsilon^n) + \frac{1}{2} [\gamma |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_1^2 + \gamma^{-1} |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_h^2 + \gamma^{-1} \tau_n (\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla W_\varepsilon^n)] \\ \leq \mathcal{J}(\Theta_\varepsilon^{n-1}, \underline{U}_\varepsilon^n), \end{aligned} \quad (2.80a)$$

where

$$\mathcal{J}(\Theta_\varepsilon^n, \underline{U}_\varepsilon^n) := \frac{1}{2} [\gamma |\Theta_\varepsilon^n|_1^2 - \gamma^{-1} |\Theta_\varepsilon^n|_h^2] + \left[\int_\Omega E(\Theta_\varepsilon^n, \underline{U}_\varepsilon^n) \, dx - \int_{\partial\Omega} \underline{g} \cdot \underline{U}_\varepsilon^n \, ds \right] \geq \mathcal{J}_0 > -\infty. \quad (2.80b)$$

Furthermore it holds that

$$\begin{aligned} \gamma^2 (G_\varepsilon(\Theta_\varepsilon^n) - G_\varepsilon(\Theta_\varepsilon^{n-1}), 1)^h + \frac{\gamma^2}{2} \tau_n |\Delta^h \Theta_\varepsilon^n|_h^2 \\ \leq \varepsilon^{-1} \gamma^2 |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_h^2 + \gamma \tau_n (\nabla W_\varepsilon^n, \nabla[\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}]) \\ + \tau_n [(\nabla \Theta_\varepsilon^n, \nabla \Theta_\varepsilon^{n-1}) + C |\underline{\mathcal{E}}(\underline{U}_\varepsilon^n)|_{0,4}^4]. \end{aligned} \quad (2.81)$$

Proof. As (2.12a) is a linear finite dimensional system, existence of $\underline{U}_\varepsilon^n$ follows from uniqueness. Given $\Theta_\varepsilon^{n-1} \in K^h$, it follows from (1.4), (1.10) and (1.21) that

$$(c(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\mathcal{E}}(\underline{U}), \underline{\mathcal{E}}(\underline{U})) \geq c_0 (\mathcal{C} \underline{\mathcal{E}}(\underline{U}), \underline{\mathcal{E}}(\underline{U})) \geq c_0 m_C |\underline{\mathcal{E}}(\underline{U})|_0^2 \geq C \|\underline{U}\|_1^2 \quad \forall \underline{U} \in \widehat{S}^h.$$

Hence we have existence and uniqueness of $\underline{U}_\varepsilon^n \in \widehat{S}^h$ solving (2.12a).

In order to prove existence of a solution $\{\Theta_\varepsilon^n, W_\varepsilon^n\} \in K^h \times S^h$ to (2.12b,c), we introduce, similarly to (2.23), for $q^h \in K^h$ the discrete anisotropic Green's operator $\mathcal{G}_{q^h}^h : Z^h \rightarrow Z^h$ such that

$$(\Xi_\varepsilon(q^h) \nabla[\mathcal{G}_{q^h}^h z^h], \nabla \chi) = (z^h, \chi)^h \quad \forall \chi \in S^h. \quad (2.82)$$

It follows immediately from (2.78) and (2.21) that $\mathcal{G}_{q^h}^h$ is well-posed. Choosing $\chi \equiv 1$ in (2.12b) yields $f\Theta_\varepsilon^n = f\Theta_\varepsilon^{n-1}$. It then follows from (2.12b) and (2.82) that

$$W_\varepsilon^n \equiv -\gamma \mathcal{G}_{\Theta_\varepsilon^{n-1}}^h \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right] + \lambda^n, \quad (2.83)$$

where $\lambda^n \in \mathbb{R}$ is a constant. Hence (2.12b,c) can be restated as: Find $\Theta_\varepsilon^n \in K^h$ and a Lagrange multiplier $\lambda^n \in \mathbb{R}$ such that

$$\begin{aligned} & \gamma (\nabla \Theta_\varepsilon^n, \nabla (\chi - \Theta_\varepsilon^n)) + (\gamma \mathcal{G}_{\Theta_\varepsilon^{n-1}}^h \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right] - \lambda^n - \gamma^{-1} \Theta_\varepsilon^{n-1}, \chi - \Theta_\varepsilon^n)^h \\ & \geq -\frac{1}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \chi - \Theta_\varepsilon^n) \quad \forall \chi \in K^h. \end{aligned} \quad (2.84)$$

It follows from (2.84) that $\Theta_\varepsilon^n \in K^h(\Theta_\varepsilon^{n-1}) := \{\chi \in K^h : \chi - \Theta_\varepsilon^{n-1} \in Z^h\}$ is such that

$$\begin{aligned} & \gamma (\nabla \Theta_\varepsilon^n, \nabla (\chi - \Theta_\varepsilon^n)) + (\gamma \mathcal{G}_{\Theta_\varepsilon^{n-1}}^h \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right] - \gamma^{-1} \Theta_\varepsilon^{n-1}, \chi - \Theta_\varepsilon^n)^h \\ & \geq -\frac{1}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \chi - \Theta_\varepsilon^n) \quad \forall \chi \in K^h(\Theta_\varepsilon^{n-1}). \end{aligned} \quad (2.85)$$

There exists a unique $\Theta_\varepsilon^n \in K^h(\Theta_\varepsilon^{n-1})$ solving (2.85) since, on noting (2.82), this is the Euler–Lagrange variational inequality of the convex minimization problem

$$\begin{aligned} \min_{z^h \in K^h(\Theta_\varepsilon^{n-1})} & \left\{ \frac{\gamma}{2} |z^h|_1^2 + \frac{\gamma}{2\tau_n} [|\Xi_\varepsilon(\Theta_\varepsilon^{n-1})|]^{1/2} \nabla \mathcal{G}_{\Theta_\varepsilon^{n-1}}^h(z^h - \Theta_\varepsilon^{n-1})|_0^2 \right. \\ & \left. - \gamma^{-1} (\Theta_\varepsilon^{n-1}, z^h)^h + \frac{1}{2} (c(z^h) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)) \right\}. \end{aligned}$$

Existence of the Lagrange multiplier λ^n in (2.84) then follows from standard optimisation theory, see e.g. Ciarlet (1988). Therefore, on noting (2.83), we have existence of a solution $\{\Theta_\varepsilon^n, W_\varepsilon^n\} \in K^h \times S^h$ to (2.12b,c). If $|\Theta_\varepsilon^n(p_j)| < 1$ for some $j \in J$ then $\pi^h[1 - (\Theta_\varepsilon^n)^2] \neq 0$ and choosing $\chi \equiv \Theta_\varepsilon^n \pm \varsigma \pi^h[1 - (\Theta_\varepsilon^n)^2]$ in (2.84) for $\varsigma > 0$ sufficiently small yields uniqueness of λ^n and, on noting (2.83), uniqueness of W_ε^n .

It follows from (1.3), (1.10), (1.13), a trace inequality and (1.21) that

$$\begin{aligned} \mathcal{J}(\Theta_\varepsilon^n, \underline{U}_\varepsilon^n) & \geq -\frac{1}{2} \gamma^{-1} \underline{m}(\Omega) + \left[\int_\Omega E(\Theta_\varepsilon^n, \underline{U}_\varepsilon^n) \, dx - \int_{\partial\Omega} \underline{g} \cdot \underline{U}_\varepsilon^n \, ds \right] \\ & \geq -\frac{1}{2} \gamma^{-1} \underline{m}(\Omega) + \frac{1}{2} c_0 m_C |\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_0^2 - \|\underline{g}\|_{0,\infty,\partial\Omega} \|\underline{U}_\varepsilon^n\|_{0,1,\partial\Omega} \\ & \geq -\frac{1}{2} \gamma^{-1} \underline{m}(\Omega) + \frac{1}{2} c_0 m_C |\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_0^2 - C |\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_0 \geq \mathcal{J}_0 > -\infty. \end{aligned} \quad (2.86)$$

Furthermore, choosing $\chi \equiv W_\varepsilon^n$ in (2.12b) and $\chi \equiv \Theta_\varepsilon^{n-1}$ in (2.12c) yields that

$$\gamma (\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}, W_\varepsilon^n)^h + \tau_n (\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla W_\varepsilon^n) = 0, \quad (2.87a)$$

$$\begin{aligned} \gamma (\nabla \Theta_\varepsilon^n, \nabla [\Theta_\varepsilon^{n-1} - \Theta_\varepsilon^n]) & \geq (W_\varepsilon^n + \gamma^{-1} \Theta_\varepsilon^{n-1}, \Theta_\varepsilon^{n-1} - \Theta_\varepsilon^n)^h \\ & \quad - \frac{1}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n), \Theta_\varepsilon^{n-1} - \Theta_\varepsilon^n). \end{aligned} \quad (2.87b)$$

On noting the fact that $c'(\Theta_\varepsilon^{n-1}) [\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}] = c(\Theta_\varepsilon^n) - c(\Theta_\varepsilon^{n-1})$, as c is affine linear, and the elementary identity

$$2r(r-s) = (r^2 - s^2) + (r-s)^2 \quad \forall r, s \in \mathbb{R}, \quad (2.88)$$

it follows from (2.87a,b), (1.3) and (2.86) that the desired result (2.80a,b) holds.

Choosing $\chi \equiv \pi^h[G'_\varepsilon(\Theta_\varepsilon^{n-1})]$ in (2.12b), and noting (2.9b) yields that

$$\gamma(\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}, G'_\varepsilon(\Theta_\varepsilon^{n-1}))^h + \tau_n(\nabla W_\varepsilon^n, \nabla \Theta_\varepsilon^{n-1}) = 0. \quad (2.89)$$

We now apply an argument similar to that in Barrett, Blowey, and Garcke (2001, Theorem 2.3). From (2.12c) we have for all $j \in J$ on choosing $\chi \equiv \Theta_\varepsilon^n + \varsigma \chi_j$, $\Theta_\varepsilon^n \pm \varsigma \chi_j$, $\Theta_\varepsilon^n - \varsigma \chi_j \in K^h$, respectively for $\varsigma > 0$ sufficiently small, that

$$\begin{aligned} \gamma(\nabla \Theta_\varepsilon^n, \nabla \chi_j) - (W_\varepsilon^n + \gamma^{-1} \Theta_\varepsilon^{n-1}, \chi_j)^h + \frac{1}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(U_\varepsilon^n), \chi_j) \\ \left\{ \begin{array}{l} \geq 0 \\ = 0 \\ \leq 0 \end{array} \right. \quad \text{if } \Theta_\varepsilon^n(p_j) \left\{ \begin{array}{l} = -1 \\ \in (-1, 1) \\ = 1 \end{array} \right. \quad . \quad (2.90) \end{aligned}$$

From (2.25), (2.2) and (2.1) it follows for all $j \in J$ that

$$\Theta_\varepsilon^n(p_j) = \pm 1 \implies \pm \Theta_\varepsilon^n(p_j) \geq \pm \Theta_\varepsilon^n(p_i) \quad \forall i \in J \implies \pm \Delta^h \Theta_\varepsilon^n(p_j) \leq 0. \quad (2.91)$$

Combining (2.90) and (2.91), and noting (2.25), (1.10), (2.3) and (2.17), yields that

$$\begin{aligned} \gamma^2 |\Delta^h \Theta_\varepsilon^n|_h^2 &= -\gamma^2 (\nabla \Theta_\varepsilon^n, \nabla (\Delta^h \Theta_\varepsilon^n)) \\ &\leq -(\gamma W_\varepsilon^n + \Theta_\varepsilon^{n-1}, \Delta^h \Theta_\varepsilon^n)^h + \frac{\gamma}{2} (c'(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(U_\varepsilon^n), \Delta^h \Theta_\varepsilon^n) \\ &\leq (\nabla[\gamma W_\varepsilon^n + \Theta_\varepsilon^{n-1}], \nabla \Theta_\varepsilon^n) + \frac{\gamma^2}{2} |\Delta^h \Theta_\varepsilon^n|_h^2 + C |\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,4}^4. \end{aligned} \quad (2.92)$$

It follows from (2.89), (2.8) and (2.92) that

$$\begin{aligned} \gamma^2 (G_\varepsilon(\Theta_\varepsilon^n) - G_\varepsilon(\Theta_\varepsilon^{n-1}), 1)^h + \frac{\gamma^2}{2} \tau_n |\Delta^h \Theta_\varepsilon^n|_h^2 \\ \leq \gamma^2 (\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}, G'_\varepsilon(\Theta_\varepsilon^n))^h + \tau_n (\nabla[\gamma W_\varepsilon^n + \Theta_\varepsilon^{n-1}], \nabla \Theta_\varepsilon^n) + C \tau_n |\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,4}^4 \\ \leq \gamma^2 (\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}, G'_\varepsilon(\Theta_\varepsilon^n) - G'_\varepsilon(\Theta_\varepsilon^{n-1}))^h + \tau_n \gamma (\nabla W_\varepsilon^n, \nabla[\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}]) \\ \quad + \tau_n (\nabla \Theta_\varepsilon^n, \nabla \Theta_\varepsilon^{n-1}) + C \tau_n |\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,4}^4 \\ \leq \varepsilon^{-1} \gamma^2 |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_h^2 + \tau_n [\gamma (\nabla W_\varepsilon^n, \nabla[\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}]) + (\nabla \Theta_\varepsilon^n, \nabla \Theta_\varepsilon^{n-1}) + C |\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,4}^4] \end{aligned}$$

and hence the desired result (2.81). \square

REMARK. 2.3 We note that (2.80a,b) and (2.81) are the discrete analogues of the energy estimates (1.16) and (1.17), respectively.

LEMMA. 2.8 *Let the assumptions of Lemmas 2.4 and 2.7 hold. Then for all $p \in [2, 2 + \delta_1]$ and for all $h \in (0, h_0)$*

$$|\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,p} \leq C. \quad (2.93)$$

Proof. Similarly to (2.76), it follows from (2.72), (2.12a), (1.13), (1.21) and a trace inequality that

$$|\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_{0,p} \leq C \sup_{\underline{0} \neq \underline{\chi} \in \widehat{\mathcal{S}}^h} \frac{|\int_{\partial\Omega} \underline{g} \cdot \underline{\chi} \, ds|}{|\underline{\underline{\mathcal{E}}}(\underline{\chi})|_{0,q}} \leq C \sup_{\underline{0} \neq \underline{\chi} \in \widehat{\mathcal{S}}^h} \frac{|\underline{\chi}|_{0,1,\partial\Omega}}{|\underline{\underline{\mathcal{E}}}(\underline{\chi})|_{0,q}} \leq C \sup_{\underline{0} \neq \underline{\chi} \in \widehat{\mathcal{S}}^h} \frac{|\underline{\chi}|_{0,1,\partial\Omega}}{\|\underline{\chi}\|_{1,q}} \leq C;$$

and hence the desired result (2.93). \square

On recalling (1.13), we set

$$\widetilde{U}_\varepsilon^n := \underline{U}_\varepsilon^n - \underline{\underline{S}}^* \underline{x} \quad (2.94)$$

as it is easier, by exploiting (2.36), to bound $|L^h \widetilde{U}_\varepsilon^n|_0$ than to bound $|L^h \underline{U}_\varepsilon^n|_0$; see the Lemma below.

LEMMA. 2.9 *Let the assumptions of Lemmas 2.2 and 2.8 hold. Assuming that $\Theta_\varepsilon^{n-1} = 1$ on $\partial\Omega$, it holds that*

$$|L^h(\widetilde{U}_\varepsilon^n)|_0 \leq C |\Theta_\varepsilon^{n-1}|_{1,4}^2. \quad (2.95)$$

Moreover, for all $h \in (0, h_0)$

$$|\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_{0,4}^4 \leq C(\delta_1) [|\Theta_\varepsilon^{n-1}|_{1,4}^{4-\delta_1} + 1]. \quad (2.96)$$

Proof. For ease of notation, let $c^{n-1} := c(\Theta_\varepsilon^{n-1})$. Assuming that $\Theta_\varepsilon^{n-1} = 1$ on $\partial\Omega$, it follows from (1.13) and $\underline{\underline{\mathcal{E}}}(\underline{\underline{S}}^* \underline{x}) = \underline{\underline{S}}^* \underline{\underline{\mathcal{E}}}(\underline{x}) = \underline{\underline{S}}^*$ that

$$\begin{aligned} \int_{\partial\Omega} \underline{g} \cdot \underline{\eta} \, ds &= \int_{\partial\Omega} c^{n-1} (\mathcal{C} \underline{\underline{S}}^* \underline{\nu}) \cdot \underline{\eta} \, ds = (\nabla \cdot (c^{n-1} \mathcal{C} \underline{\underline{S}}^* \underline{\eta}), 1) \\ &= (c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{\underline{S}}^* \underline{x}), \underline{\underline{\mathcal{E}}}(\underline{\eta})) + (\nabla \cdot (c^{n-1} \mathcal{C} \underline{\underline{S}}^*), \underline{\eta}) \quad \forall \underline{\eta} \in \underline{\underline{H}}^1(\Omega). \end{aligned} \quad (2.97)$$

Combining (2.12a), (2.94), and (2.97) yields that

$$(c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\widetilde{U}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\underline{\chi})) = (\nabla \cdot (c^{n-1} \underline{\underline{S}}), \underline{\chi}) \quad \forall \underline{\chi} \in \underline{\underline{S}}^h. \quad (2.98)$$

For the ensuing analysis it is convenient to introduce $\widetilde{u}_\varepsilon^n \in \widehat{\underline{\underline{H}}^1}(\Omega)$ such that

$$(c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\widetilde{u}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\underline{\eta})) = (\nabla \cdot (c^{n-1} \underline{\underline{S}}), \underline{\eta}) \quad \forall \underline{\eta} \in \underline{\underline{H}}^1(\Omega). \quad (2.99)$$

Existence and uniqueness of $\widetilde{u}_\varepsilon^n$, and the bound

$$\|\widetilde{u}_\varepsilon^n\|_1 \leq C \quad (2.100)$$

are easily established on noting (2.97), (1.13), (1.4), (1.10), (1.21) and a trace inequality.

We now address the $H^2(\Omega)$ regularity of $\widetilde{u}_\varepsilon^n$. If $\widehat{\underline{\eta}} \in \underline{\underline{H}}^1(\Omega)$, then $\underline{\eta} := [c^{n-1}]^{-1} \widehat{\underline{\eta}}$ satisfies, on noting (1.4) and (1.18),

$$|\underline{\eta}|_1 \leq C [|\widehat{\underline{\eta}}|_1 + |\Theta_\varepsilon^{n-1}|_{1,2+\varsigma} |\widehat{\underline{\eta}}|_{0, \frac{2(2+\varsigma)}{\varsigma}}] \leq C [1 + |\Theta_\varepsilon^{n-1}|_{1,2+\varsigma}] \|\widehat{\underline{\eta}}\|_1, \quad \varsigma > 0,$$

and hence $\underline{\eta} \in \underline{H}^1(\Omega)$. Choosing $\underline{\eta} \equiv [c^{n-1}]^{-1} \widehat{\underline{\eta}}$ in (2.99) yields, on noting (2.54), that for all $\widehat{\underline{\eta}} \in \underline{H}^1(\Omega)$

$$(\mathcal{C} \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\widehat{\underline{\eta}})) = ([c^{n-1}]^{-1} \nabla \cdot (c^{n-1} \underline{\underline{\mathcal{S}}}) - c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n) \nabla [c^{n-1}]^{-1}, \widehat{\underline{\eta}}). \quad (2.101)$$

It follows from (2.101), (2.34), (1.18), (2.100) and (1.4) that

$$\begin{aligned} \|\tilde{\underline{u}}_\varepsilon^n\|_2 &\leq C [|\underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n)|_{0,4} |c^{n-1}|_{1,4} + |c^{n-1}|_1] \leq C [|\tilde{\underline{u}}_\varepsilon^n|_1^{\frac{1}{2}} \|\tilde{\underline{u}}_\varepsilon^n\|_2^{\frac{1}{2}} + 1] |c^{n-1}|_{1,4} \\ &\leq C |c^{n-1}|_{1,4}^2 \leq C |\Theta_\varepsilon^{n-1}|_{1,4}^2. \end{aligned} \quad (2.102)$$

From (1.10), (1.4), (1.21), (2.99), (2.98), (1.9), (2.15) and (2.102) we have that

$$\begin{aligned} C_1 \|\tilde{\underline{u}}_\varepsilon^n - \tilde{\underline{U}}_\varepsilon^n\|_1^2 &\leq (c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n - \tilde{\underline{U}}_\varepsilon^n), \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n - \tilde{\underline{U}}_\varepsilon^n)) \\ &\leq (c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}((I - \pi^h)\tilde{\underline{u}}_\varepsilon^n), \underline{\underline{\mathcal{E}}}((I - \pi^h)\tilde{\underline{u}}_\varepsilon^n)) \leq C_2 |(I - \pi^h)\tilde{\underline{u}}_\varepsilon^n|_1^2 \leq C_3 h^2 |\tilde{\underline{u}}_\varepsilon^n|_2^2. \end{aligned} \quad (2.103)$$

It follows from (2.35) and (2.103) that

$$\begin{aligned} |L^h \tilde{\underline{U}}_\varepsilon^n|_0 &\leq |L^h(\tilde{\underline{U}}_\varepsilon^n - \pi^h \tilde{\underline{u}}_\varepsilon^n)|_0 + |L^h(\pi^h \tilde{\underline{u}}_\varepsilon^n)|_0 \leq C h^{-1} |\tilde{\underline{U}}_\varepsilon^n - \pi^h \tilde{\underline{u}}_\varepsilon^n|_1 + |L^h(\pi^h \tilde{\underline{u}}_\varepsilon^n)|_0 \\ &\leq C h^{-1} [|\tilde{\underline{U}}_\varepsilon^n - \tilde{\underline{u}}_\varepsilon^n|_1 + |(I - \pi^h)\tilde{\underline{u}}_\varepsilon^n|_1] + |L^h(\pi^h \tilde{\underline{u}}_\varepsilon^n)|_0 \leq C |\tilde{\underline{u}}_\varepsilon^n|_2 + |L^h(\pi^h \tilde{\underline{u}}_\varepsilon^n)|_0. \end{aligned} \quad (2.104)$$

It follows from (2.99) and (1.4) that

$$c^{n-1} \mathcal{C} \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n) \underline{\nu} = \underline{0} \quad \text{on } \partial\Omega \quad \implies \quad \mathcal{C} \underline{\underline{\mathcal{E}}}(\tilde{\underline{u}}_\varepsilon^n) \underline{\nu} = \underline{0} \quad \text{on } \partial\Omega. \quad (2.105)$$

The desired result (2.95) then follows from (2.104), (2.105), (2.36) and (2.102).

It follows from (2.94) that

$$|\underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n)|_{0,4}^4 \leq C [|\underline{\underline{\mathcal{E}}}(\tilde{\underline{U}}_\varepsilon^n)|_{0,4}^4 + 1]. \quad (2.106)$$

On noting (2.35), we have for any $\alpha \in (0, 1)$ that

$$|\underline{\underline{\mathcal{E}}}(\tilde{\underline{U}}_\varepsilon^n)|_{0,4}^4 \leq C(\alpha) |\underline{\underline{\mathcal{E}}}(\tilde{\underline{U}}_\varepsilon^n)|_{0,2+2\alpha}^{2+\alpha} |\underline{\underline{\mathcal{E}}}(\tilde{\underline{U}}_\varepsilon^n)|_{0,\frac{(2-\alpha)(2+2\alpha)}{\alpha}}^{2-\alpha} \leq C(\alpha) |\underline{\underline{\mathcal{E}}}(\tilde{\underline{U}}_\varepsilon^n)|_{0,2+2\alpha}^{2+\alpha} |L^h \tilde{\underline{U}}_\varepsilon^n|_0^{2-\alpha}. \quad (2.107)$$

Combining (2.106), (2.107), (2.94), (2.93) and (2.95) yields the desired result (2.96). \square

LEMMA. 2.10 *Let $\theta^0 \in K \cap H^2(\Omega)$ with $\frac{\partial \theta^0}{\partial \underline{\nu}} = 0$ on $\partial\Omega$, and the assumptions (A) hold. On choosing $\Theta_\varepsilon^0 \equiv \pi^h \theta^0$ it follows that $\Theta_\varepsilon^0 \in K^h$ is such that for all $h > 0$*

$$\|\Theta_\varepsilon^0\|_1^2 + |\Delta^h \Theta_\varepsilon^0|_h^2 + (G_\varepsilon(\Theta_\varepsilon^0), 1)^h \leq C. \quad (2.108)$$

Proof. The desired result (2.108) follows immediately from (2.15), (2.28), (2.3), (2.17), (2.7) and (2.6). \square

THEOREM. 2.1 *Let the assumptions of Lemma 2.10 hold. Then for all $\varepsilon \in (0, 1)$, $h \in (0, h_0)$ and for all time partitions $\{\tau_n\}_{n=1}^N$, the solution $\{\underline{U}_\varepsilon^n, \Theta_\varepsilon^n, W_\varepsilon^n\}_{n=1}^N$ to $(P_\varepsilon^{h,\tau})$ is such that $f\Theta_\varepsilon^n = f\Theta_\varepsilon^0$, $n = 1 \rightarrow N$, and*

$$\begin{aligned} & \gamma \max_{n=1 \rightarrow N} \|\Theta_\varepsilon^n\|_1^2 + \sum_{n=1}^N [\gamma |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_1^2 + \gamma^{-1} (|\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_0^2 + \tau_n (\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla W_\varepsilon^n))] \\ & + \sum_{n=2}^N (c(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1}), \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1})) \leq C [1 + \|\Theta_\varepsilon^0\|_1^2] \leq C. \end{aligned} \quad (2.109)$$

In addition

$$\gamma \sum_{n=1}^N \tau_n \left| \mathcal{G} \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right] \right|_1^2 + \gamma \tau^{-\frac{1}{2}} \sum_{n=1}^N |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_0^2 \leq C [1 + \|\Theta_\varepsilon^0\|_1^2] \leq C. \quad (2.110)$$

Moreover, on assuming (2.34) holds, if \mathcal{C} is anisotropic, $\tau_n \leq C \tau_{n-1}$, $n = 2 \rightarrow N$, and $\Theta_\varepsilon^{n-1} = 1$ on $\partial\Omega$, $n = 1 \rightarrow N$; then

$$\gamma^2 \max_{n=1 \rightarrow N} (G_\varepsilon(\Theta_\varepsilon^n), 1)^h + \sum_{n=1}^N \tau_n [\gamma^2 |\Delta^h \Theta_\varepsilon^n|_h^2 + \|\underline{U}_\varepsilon^n\|_{1,4}^4] \leq C(T) [1 + \varepsilon^{-1} \tau^{\frac{1}{2}}]. \quad (2.111)$$

Proof. First, it follows from (2.80b), (1.3), (2.88) and (2.12a) that for $n = 2 \rightarrow N$

$$\mathcal{J}(\Theta_\varepsilon^{n-1}, \underline{U}_\varepsilon^n) = \mathcal{J}(\Theta_\varepsilon^{n-1}, \underline{U}_\varepsilon^{n-1}) - \frac{1}{2} (c(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1}), \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1})). \quad (2.112)$$

Summing (2.80a) from $n = 1 \rightarrow k$ and noting (2.112), (2.80b), (1.3), (1.10), (1.4) and a trace inequality, yields for $k = 2 \rightarrow N$ that

$$\begin{aligned} & \mathcal{J}(\Theta_\varepsilon^k, \underline{U}_\varepsilon^k) + \frac{1}{2} \sum_{n=1}^k [\gamma |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_1^2 + \gamma^{-1} (|\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_h^2 + \tau_n (\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla W_\varepsilon^n))] \\ & + \frac{1}{2} \sum_{n=2}^k (c(\Theta_\varepsilon^{n-1}) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1}), \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^n - \underline{U}_\varepsilon^{n-1})) \\ & \leq \mathcal{J}(\Theta_\varepsilon^0, \underline{U}_\varepsilon^1) \leq C [1 + \|\Theta_\varepsilon^0\|_1 + \|\underline{U}_\varepsilon^1\|_1^2]. \end{aligned} \quad (2.113)$$

The desired result (2.109) then follows from (2.80a) for $n = 1$, (2.113) for $k = 2 \rightarrow N$, (2.80b), (2.3), (2.17) and the fact that $\Theta_\varepsilon^n \in K^h$, $n = 0 \rightarrow N$, (2.93), (1.21) and (2.108).

From (2.20), (2.4), (2.12b), (2.78) and (2.19) we obtain for any $\eta \in H^1(\Omega)$ that

$$\begin{aligned} \gamma \left(\nabla \mathcal{G} \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right], \nabla \eta \right) &= \gamma \left(\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n}, \eta \right) = \gamma \left(\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n}, Q^h \eta \right)^h \\ &= -(\Xi_\varepsilon(\Theta_\varepsilon^{n-1}) \nabla W_\varepsilon^n, \nabla [Q^h \eta]) \leq \|[\Xi_\varepsilon(\Theta_\varepsilon^{n-1})]\|_0^{\frac{1}{2}} \|\nabla W_\varepsilon^n\|_0 \|Q^h \eta\|_1 \\ &\leq C \|[\Xi_\varepsilon(\Theta_\varepsilon^{n-1})]\|_0^{\frac{1}{2}} \|\nabla W_\varepsilon^n\|_0 \|\eta\|_1. \end{aligned} \quad (2.114)$$

The first bound in (2.110) then follows from (2.114) and (2.109). Moreover, we have from (2.20) that

$$\sum_{n=1}^N |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_0^2 \leq \tau^{\frac{1}{2}} \left[\sum_{n=1}^N |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_1^2 \right]^{\frac{1}{2}} \left[\sum_{n=1}^N \tau_n \left| \mathcal{G} \left[\frac{\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}}{\tau_n} \right] \right|_1^2 \right]^{\frac{1}{2}}.$$

The second bound in (2.110) then follows from the first and (2.109).

Finally, summing (2.81) from $n = 1 \rightarrow k$ and noting (2.3), (2.17), (2.78), (2.109), (2.110), (2.108), (2.96), (2.27b), our assumption on τ_n , and (1.25) yields for any $k \leq N$ that

$$\begin{aligned} & \gamma^2 (G_\varepsilon(\Theta_\varepsilon^k), 1)^h + \frac{\gamma^2}{2} \sum_{n=1}^k \tau_n |\Delta^h \Theta_\varepsilon^n|_h^2 \\ & \leq \gamma^2 (G_\varepsilon(\Theta_\varepsilon^0), 1)^h + 4\varepsilon^{-1} \gamma^2 \sum_{n=1}^k |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_0^2 + t_k \max_{n=0 \rightarrow k} \|\Theta_\varepsilon^n\|_1^2 + C \sum_{n=1}^k \tau_n |\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)|_{0,4}^4 \\ & \quad + \left[\varepsilon^{-1} \sum_{n=1}^k \tau_n |[\Xi_\varepsilon(\Theta_\varepsilon^{n-1})]^{\frac{1}{2}} \nabla W_\varepsilon^n|_0^2 \right]^{\frac{1}{2}} \left[\gamma \sum_{n=1}^k \tau_n |\Theta_\varepsilon^n - \Theta_\varepsilon^{n-1}|_1^2 \right]^{\frac{1}{2}} \\ & \leq C(T) [1 + \varepsilon^{-1} \tau^{\frac{1}{2}}] + C \sum_{n=1}^k \tau_n |\Theta_\varepsilon^{n-1}|_{1,4}^{4-\delta_1} \\ & \leq C(T) [1 + \varepsilon^{-1} \tau^{\frac{1}{2}}] + C \sum_{n=2}^k \tau_{n-1} |\Delta^h \Theta_\varepsilon^{n-1}|_h^{2-\frac{\delta_1}{2}} \leq C(T) [1 + \varepsilon^{-1} \tau^{\frac{1}{2}}]. \end{aligned} \quad (2.115)$$

Hence the desired result (2.111) follows immediately from (2.115) and (1.21). \square

REMARK. 2.4 The approximation $(\mathbb{P}_\varepsilon^{h,\tau})$ of (P) requires solving for $\{\Theta_\varepsilon^n, W_\varepsilon^n\}$ over the whole domain Ω , due to the non-degeneracy of $\Xi_\varepsilon(\cdot)$, see (2.78). For computational speed it would be more convenient to solve for $\{\Theta_\varepsilon^n, W_\varepsilon^n\}$ just in the interfacial region, $|\Theta_\varepsilon^{n-1}| < 1$. With this in mind, and adopting the notation (2.10) and (2.11), we introduce $\Xi_\varepsilon^* : S^h \rightarrow [L^\infty(\Omega)]^{2 \times 2}$ such that $\Xi_\varepsilon^*(z^h)|_\sigma := R_\sigma \widehat{\Xi}_\varepsilon^*(\widehat{z}^h)|_\sigma R_\sigma^T$, where

$$[\widehat{\Xi}_\varepsilon^*(\widehat{z}^h)|_\sigma]_{kk} := \begin{cases} 0 & \text{if } \widehat{z}^h(\widehat{p}_k) = \widehat{z}^h(\widehat{p}_0) = \pm 1, \\ [\widehat{\Xi}_\varepsilon(\widehat{z}^h)|_\sigma]_{kk} & \text{otherwise.} \end{cases}$$

We note that the key identities, $\Xi_\varepsilon(z^h)$ in (2.9a,b) replaced by $\Xi_\varepsilon^*(z^h)$, still hold. We then introduce the approximation $(\widetilde{\mathbb{P}}_\varepsilon^{h,\tau})$ of (P), which is the same as $(\mathbb{P}_\varepsilon^{h,\tau})$ but with $\Xi_\varepsilon(\Theta_\varepsilon^{n-1})$ in (2.12b) replaced by $\Xi_\varepsilon^*(\Theta_\varepsilon^{n-1})$. As $\Xi_\varepsilon^*(\cdot)$ is now degenerate, existence of a solution $\{\Theta_\varepsilon^n, W_\varepsilon^n\}$ to $(\widetilde{\mathbb{P}}_\varepsilon^{h,\tau})$ does not appear to be trivial. However, this can easily be established by splitting the nodes into passive and active sets, see e.g. Barrett, Blowey, and Garcke (1999). Moreover, one can show that Θ_ε^n is unique, and $W_\varepsilon^n(p_j)$ is unique if $(\Xi_\varepsilon^*(\Theta_\varepsilon^{n-1}), \chi_j) > 0$. Furthermore, one can establish analogues of the energy estimates (2.109) and (2.110). Unfortunately, it does not appear possible to establish an analogue of the key energy estimate (2.111) for $(\widetilde{\mathbb{P}}_\varepsilon^{h,\tau})$.

3 Convergence

In this section we will show convergence of the discrete solutions obtained in Section 2 to a weak solution of problem (P). We will use methods developed by Barrett, Blowey, and Garcke (1998), Grün (2003) and Barrett, Nürnberg, and Styles (2004) to deal with the degeneracy of b . Furthermore, it will be crucial to show strong convergence of $\underline{\underline{\mathcal{E}}}(U_\varepsilon^n)$ in order to pass to the limit in the nonlinearity $\mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^n) : \underline{\underline{\mathcal{E}}}(U_\varepsilon^n)$.

Let

$$\Theta_\varepsilon(\cdot, t) := \frac{t-t_{n-1}}{\tau_n} \Theta_\varepsilon^n(\cdot) + \frac{t_n-t}{\tau_n} \Theta_\varepsilon^{n-1}(\cdot) \quad t \in [t_{n-1}, t_n] \quad n \geq 1, \quad (3.1a)$$

$$\Theta_\varepsilon^+(\cdot, t) := \Theta_\varepsilon^n(\cdot), \quad \Theta_\varepsilon^-(\cdot, t) := \Theta_\varepsilon^{n-1}(\cdot) \quad t \in (t_{n-1}, t_n] \quad n \geq 1. \quad (3.1b)$$

We note for future reference that

$$\Theta_\varepsilon(\cdot, t) - \Theta_\varepsilon^\pm(\cdot, t) = (t - t_n^\pm) \frac{\partial \Theta_\varepsilon}{\partial t}(\cdot, t) \quad t \in (t_{n-1}, t_n) \quad n \geq 1, \quad (3.2)$$

where $t_n^+ := t_n$ and $t_n^- := t_{n-1}$. We introduce also

$$\bar{\tau}(t) := \tau_n \quad t \in (t_{n-1}, t_n] \quad n \geq 1. \quad (3.3)$$

Using the above notation, and introducing analogous notation for W_ε and $\underline{U}_\varepsilon$, $(P_\varepsilon^{h,\tau})$ can be restated as: Find $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\} \in L^\infty(0, T; \widehat{S}^h) \times C([0, T]; K^h) \times L^\infty(0, T; S^h)$ such that

$$\int_0^T (c(\Theta_\varepsilon^-) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^+), \underline{\underline{\mathcal{E}}}(\chi)) dt = \int_0^T \int_{\partial\Omega} \underline{g} \cdot \underline{\chi} ds dt \quad \forall \chi \in L^\infty(0, T; \underline{S}^h), \quad (3.4a)$$

$$\int_0^T \left[\gamma \left(\frac{\partial \Theta_\varepsilon}{\partial t}, \chi \right)^h + (\Xi_\varepsilon(\Theta_\varepsilon^-) \nabla W_\varepsilon^+, \nabla \chi) \right] dt = 0 \quad \forall \chi \in L^\infty(0, T; S^h), \quad (3.4b)$$

$$\begin{aligned} \gamma \int_0^T (\nabla \Theta_\varepsilon^+, \nabla [\chi - \Theta_\varepsilon^+]) dt &\geq \int_0^T (W_\varepsilon^+ + \gamma^{-1} \Theta_\varepsilon^-, \chi - \Theta_\varepsilon^+)^h dt \\ &\quad - \frac{1}{2} \int_0^T (c'(\Theta_\varepsilon^-) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^+) : \underline{\underline{\mathcal{E}}}(U_\varepsilon^+), \chi - \Theta_\varepsilon^+) dt \quad \forall \chi \in L^\infty(0, T; K^h). \end{aligned} \quad (3.4c)$$

LEMMA. 3.1 *Let $\theta^0 \in K \cap H^2(\Omega)$ with $\frac{\partial \theta^0}{\partial \nu} = 0$ on $\partial\Omega$ and $f\theta^0 \in (-1, 1)$. Let $\{\mathcal{T}^h, \varepsilon, \{\tau_n\}_{n=1}^N\}_{h>0}$ be such that Ω and $\{\mathcal{T}^h\}_{h>0}$ fulfil assumptions (A), $\varepsilon \in (0, 1)$ with $\varepsilon \rightarrow 0$ as $h \rightarrow 0$ and $\tau_n \leq C \tau_{n-1} \leq C \varepsilon^2$, $n = 2 \rightarrow N$. Let $\Theta_\varepsilon^0 \equiv \pi^h \theta^0$. Then there exists a subsequence of $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$, where $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}$ solve $(P_\varepsilon^{h,\tau})$, and functions*

$$\theta \in L^\infty(0, T; K) \cap H^1(0, T; (H^1(\Omega))') \quad \text{and} \quad \underline{u} \in L^\infty(0, T; \widehat{\underline{V}}_{2+\delta_1}), \quad \delta_1 > 0, \quad (3.5)$$

with $\theta(\cdot, 0) = \theta^0(\cdot)$ in $L^2(\Omega)$ and $f\theta(\cdot, t) = f\theta^0$ for a.a. $t \in (0, T)$, such that as $h \rightarrow 0$

$$\Theta_\varepsilon, \Theta_\varepsilon^\pm \rightarrow \theta \quad \text{weak-* in } L^\infty(0, T; H^1(\Omega)), \quad (3.6a)$$

$$\mathcal{G} \frac{\partial \Theta_\varepsilon}{\partial t} \rightarrow \mathcal{G} \frac{\partial \theta}{\partial t} \quad \text{weakly in } L^2(0, T; H^1(\Omega)), \quad (3.6b)$$

$$\underline{U}_\varepsilon^+ \rightarrow \underline{u} \quad \text{weak-* in } L^\infty(0, T; \underline{W}^{1,2+\delta_1}(\Omega)), \quad (3.6c)$$

$$\Theta_\varepsilon, \Theta_\varepsilon^\pm \rightarrow \theta \quad \text{strongly in } L^2(0, T; L^s(\Omega)), \quad (3.7a)$$

$$\Xi_\varepsilon(\Theta_\varepsilon^-) \rightarrow b(\theta) \mathcal{I} \quad \text{strongly in } L^2(0, T; L^s(\Omega)), \quad (3.7b)$$

$$c(\Theta_\varepsilon^-) \rightarrow c(\theta) \quad \text{strongly in } L^2(0, T; L^s(\Omega)), \quad (3.7c)$$

$$\underline{U}_\varepsilon^+ \rightarrow \underline{u} \quad \text{strongly in } L^2(0, T; \underline{H}^1(\Omega)); \quad (3.7d)$$

for all $s \in [2, \infty)$. Moreover, $\{\underline{u}, \theta\}$ satisfy

$$\int_{\Omega_T} c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) : \underline{\underline{\mathcal{E}}}(\underline{\eta}) \, dx \, dt = \int_0^T \int_{\partial\Omega} \underline{g} \cdot \underline{\eta} \, ds \, dt \quad \forall \underline{\eta} \in L^2(0, T; \underline{H}^1(\Omega)). \quad (3.8)$$

Furthermore, on assuming (2.34) holds, if \mathcal{C} is anisotropic, and if

$$\Theta_\varepsilon = 1 \text{ on } \partial\Omega; \quad (3.9)$$

then $\{\theta, \underline{u}\}$, in addition to (3.5), satisfy

$$\theta \in L^2(0, T; H^2(\Omega)) \quad \text{and} \quad \underline{u} \in L^4(0, T; \underline{W}^{1,4}(\Omega)); \quad (3.10)$$

and there exists a subsequence of $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$ satisfying (3.6a–c), (3.7a–d) and as $h \rightarrow 0$

$$\Delta^h \Theta_\varepsilon, \Delta^h \Theta_\varepsilon^\pm \rightarrow \Delta \theta \quad \text{weakly in } L^2(\Omega_T), \quad (3.11a)$$

$$\Theta_\varepsilon, \Theta_\varepsilon^\pm \rightarrow \theta \quad \text{weakly in } L^2(0, T; W^{1,s}(\Omega)), \quad \text{for any } s \in [2, \infty), \quad (3.11b)$$

$$\Theta_\varepsilon \rightarrow \theta \quad \text{strongly in } L^2(0, T; C^{0,\beta}(\overline{\Omega})), \quad \text{for any } \beta \in (0, 1), \quad (3.11c)$$

$$\underline{U}_\varepsilon^+ \rightarrow \underline{u} \quad \text{weakly in } L^4(0, T; \underline{W}^{1,4}(\Omega)). \quad (3.11d)$$

Proof. On noting (3.1a,b), (3.3) and (1.21); the bounds (2.93), (2.109) and (2.110) imply that

$$\begin{aligned} & \|\underline{U}_\varepsilon^+\|_{L^\infty(0,T;\underline{W}^{1,2+\delta_1}(\Omega))}^2 + \|\Theta_\varepsilon^{(\pm)}\|_{L^\infty(0,T;H^1(\Omega))}^2 + \|[\Xi_\varepsilon(\Theta_\varepsilon^-)]^{\frac{1}{2}} \nabla W_\varepsilon^+\|_{L^2(0,T;\underline{L}^2(\Omega))}^2 \\ & + \|\bar{\tau}^{\frac{1}{2}} \frac{\partial \Theta_\varepsilon}{\partial t}\|_{L^2(0,T;H^1(\Omega))}^2 + \|\mathcal{G} \frac{\partial \Theta_\varepsilon}{\partial t}\|_{L^2(0,T;H^1(\Omega))}^2 + \tau^{-\frac{1}{2}} \|\bar{\tau}^{\frac{1}{2}} \frac{\partial \Theta_\varepsilon}{\partial t}\|_{L^2(\Omega_T)}^2 \leq C. \end{aligned} \quad (3.12)$$

Furthermore, we deduce from (3.2) and (3.12) that

$$\|\Theta_\varepsilon - \Theta_\varepsilon^\pm\|_{L^2(0,T;H^1(\Omega))}^2 \leq \|\bar{\tau} \frac{\partial \Theta_\varepsilon}{\partial t}\|_{L^2(0,T;H^1(\Omega))}^2 \leq C \tau. \quad (3.13)$$

Hence on noting (3.12), (3.13), $\Theta_\varepsilon(\cdot, t) \in K^h$, $\underline{U}_\varepsilon^+(\cdot, t) \in \widehat{S}^h$, and (1.19a) we can choose a subsequence $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$ such that the convergence results (3.5), (3.6a–c) and (3.7a)

hold. Then (3.5) and Theorem 2.1 yield, on noting (1.19b), our assumption on Θ_ε^0 and (2.16) that the subsequence satisfies the additional initial and integral conditions.

The desired results (3.7b,c) follow from (2.79), (2.14), (3.12), (2.77), (1.7) and (1.4); see Barrett, Nürnberg, and Styles (2004, Lemma 3.1) for details.

For any $\underline{\eta} \in L^2(0, T; \underline{H}^2(\Omega))$, we choose $\underline{\chi} \equiv \pi^h \underline{\eta}$ in (2.12a). The desired result (3.8) then follows from (2.15), a trace inequality, (3.12), (1.4), (3.7c), (3.6c) and a density result. We have from (3.4a) and (3.8) that

$$\begin{aligned} & \int_{\Omega_T} c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u} - \underline{U}_\varepsilon^+) : \underline{\underline{\mathcal{E}}}(\underline{u} - \underline{U}_\varepsilon^+) \, dx \, dt \\ &= \int_{\Omega_T} [c(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u} - \underline{U}_\varepsilon^+) : \underline{\underline{\mathcal{E}}}(\underline{u}) + [c(\theta) - c(\Theta_\varepsilon^-)] \mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^+) : \underline{\underline{\mathcal{E}}}(\underline{U}_\varepsilon^+)] \, dx \, dt. \end{aligned} \quad (3.14)$$

The desired result (3.7d) then follows from (3.14), on noting (1.4), (1.21), (3.6c) and (3.7c).

It follows from (2.111), (2.108), (2.3), (2.17), (3.1a,b) and our assumptions on $\{\tau_n\}_{n=1}^N$ and ε that

$$\|\Delta^h \Theta_\varepsilon^{(\pm)}\|_{L^2(\Omega_T)}^2 + \|\underline{U}_\varepsilon^+\|_{L^4(0, T; \underline{W}^{1,4}(\Omega))}^4 \leq C(T). \quad (3.15)$$

The desired results (3.10) and (3.11a,d) then follow from (3.15), (2.25), (2.16), (2.18), (3.12), (3.6a), elliptic regularity as Ω is convex polygonal, and (3.5); see Barrett, Nürnberg, and Styles (2004, Lemma 3.1) for details. Furthermore, it follows from (3.11a) and (2.27a) that (3.11b) holds on extracting a further subsequence. Finally, (3.11c) follows from (3.11b), (3.6b), (1.19a) and the compact embedding $W^{1,s}(\Omega) \hookrightarrow C^{0,\beta}(\overline{\Omega})$. \square

REMARK. 3.1 The condition $\theta^0 \in H^2(\Omega)$ with $\frac{\partial \theta^0}{\partial \nu} = 0$ can be relaxed, but it is not particularly restrictive. See e.g. Barrett and Nürnberg (2004).

In addition to the above lemma, we need the following two lemmas in order to prove our main result, Theorem 3.1 below.

LEMMA. 3.2 *Let all the assumptions of Lemma 3.1 hold. If in addition $\tau_n = \tau$, $n = 1 \rightarrow N$, then*

$$\int_0^{T-\varsigma} |\Theta_\varepsilon^\pm(t+\varsigma) - \Theta_\varepsilon^\pm(t)|_0^2 \, dt \leq C \varsigma \quad \forall \varsigma \in (0, T).$$

Moreover, it holds that the subsequence of $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$ in Lemma 3.1 is such that for any $\beta \in (0, 1)$

$$\Theta_\varepsilon^\pm \rightarrow \theta \quad \text{strongly in } L^2(0, T; C^{0,\beta}(\overline{\Omega})) \quad \text{as } h \rightarrow 0; \quad (3.16a)$$

and, on extracting a further subsequence, it holds for a.a. $t \in (0, T)$ that

$$\Theta_\varepsilon^\pm(\cdot, t) \rightarrow \theta(\cdot, t) \quad \text{strongly in } C^{0,\beta}(\overline{\Omega}) \quad \text{as } h \rightarrow 0. \quad (3.16b)$$

Proof. The proof is the same as that of Lemma 3.3 (with $\alpha = 0$) in Barrett, Nürnberg, and Styles (2004). \square

From (3.12), (2.78), (2.77), (1.7) and (3.16b) we see that we can only control ∇W_ε^+ on the set where $\Xi_\varepsilon(\Theta_\varepsilon^-)$ is bounded below independently of ε , and hence h , as $\varepsilon \rightarrow 0$ and $h \rightarrow 0$, i.e. on the set where $|\theta| < 1$. Therefore in order to construct the appropriate limits as $h \rightarrow 0$, we introduce the following open subsets of $\overline{\Omega}$. For any $\rho \in (0, 1)$, we define for *a.a.* $t \in (0, T)$

$$B_\rho(t) := \{x \in \overline{\Omega} : |\theta(x, t)| < 1 - \rho\}. \quad (3.17)$$

We have from (3.16b), see Barrett, Nürnberg, and Styles (2004) for details, that for *a.a.* $t \in (0, T)$ and any $\rho \in (0, \rho_0)$, there exists an $h_0(\rho, t)$ such that for all $h \leq h_0(\rho, t)$ there exist collections of simplices $\mathcal{T}_{B, \rho}^h(t) \subset \mathcal{T}^h$ such that

$$B_\rho(t) \subset B_\rho^h(t) := \cup_{\sigma \in \mathcal{T}_{B, \rho}^h(t)} \overline{\sigma} \subset B_{\frac{\rho}{2}}(t). \quad (3.18)$$

In addition for *a.a.* $t \in (0, T)$ and any fixed $\rho \in (0, \widehat{\rho}_0)$, where $\widehat{\rho}_0 := \min\{\rho_0, \frac{1}{2}\}$, it follows from (3.17), (3.16b) and our assumption on ε in Lemma 3.1 that there exists an $\widehat{h}_0(\rho, t) \leq h_0(\rho, t)$ such that for $h \leq \widehat{h}_0(\rho, t)$

$$1 - 2\rho \leq |\Theta_\varepsilon^\pm(x, t)| \quad \forall x \notin B_\rho(t), \quad |\Theta_\varepsilon^\pm(x, t)| \leq 1 - \frac{\rho}{2} \quad \forall x \in B_\rho(t) \quad \text{and} \quad \varepsilon \leq \rho. \quad (3.19)$$

LEMMA. 3.3 *Let all the assumptions of Lemma 3.2 hold. Then for *a.a.* $t \in (0, T)$ there exists a function*

$$w(\cdot, t) \equiv -\gamma \Delta \theta(\cdot, t) - \gamma^{-1} \theta(\cdot, t) + \frac{1}{2} [c'(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(u) : \underline{\underline{\mathcal{E}}}(u)](\cdot, t) \in H_{loc}^1(\{|\theta(\cdot, t)| < 1\}); \quad (3.20)$$

where $\{|\theta(\cdot, t)| < 1\} := \{x \in \Omega : |\theta(x, t)| < 1\}$. Moreover, on extracting a further subsequence from the subsequence $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$ in Lemma 3.2, it holds as $h \rightarrow 0$ that

$$\Xi_\varepsilon(\Theta_\varepsilon^-) \nabla W_\varepsilon^+ \rightarrow \mathcal{H}_{\{|\theta| < 1\}} b(\theta) \nabla w \quad \text{weakly in } L^2(0, T; \underline{L}^2(\Omega)); \quad (3.21)$$

where $\mathcal{H}_{\{|\theta| < 1\}}$ is the characteristic function of the set $\{|\theta| < 1\} := \{(x, t) \in \Omega_T : |\theta(x, t)| < 1\}$.

Proof. It follows from (3.12) and (2.78) that

$$\|\Xi_\varepsilon(\Theta_\varepsilon^-) \nabla W_\varepsilon^+\|_{L^2(0, T; \underline{L}^2(\Omega))}^2 \leq C. \quad (3.22)$$

Hence (3.22) implies that there exists a function $\underline{z} \in L^2(0, T; \underline{L}^2(\Omega))$, and on extracting a further subsequence from the subsequence $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$ in Lemma 3.2, it holds as $h \rightarrow 0$ that

$$\Xi_\varepsilon(\Theta_\varepsilon^-) \nabla W_\varepsilon^+ \rightarrow \underline{z} \quad \text{weakly in } L^2(0, T; \underline{L}^2(\Omega)). \quad (3.23)$$

We now identify \underline{z} .

First, we consider a fixed $\rho \in (0, \rho_0)$. It follows from (1.7), (2.77), (2.78), (3.19) and (3.12) that for *a.a.* $t \in (0, T)$ and for all $h \leq \widehat{h}_0(\rho, t)$

$$\begin{aligned} \rho \left(1 - \frac{\rho}{4}\right) |\nabla W_\varepsilon^+(\cdot, t)|_{0, B_\rho(t)}^2 &= b \left(1 - \frac{\rho}{2}\right) |\nabla W_\varepsilon^+(\cdot, t)|_{0, B_\rho(t)}^2 \leq b_\varepsilon \left(1 - \frac{\rho}{2}\right) |\nabla W_\varepsilon^+(\cdot, t)|_{0, B_\rho(t)}^2 \\ &\leq |([\Xi_\varepsilon(\Theta_\varepsilon^-)]^{\frac{1}{2}} \nabla W_\varepsilon^+(\cdot, t))|_0^2 \leq C(t). \end{aligned} \quad (3.24)$$

From (3.24), (3.18), (2.78) and (3.19) we have for *a.a.* $t \in (0, T)$ and for all $h \leq \widehat{h}_0(\rho, t)$

$$\begin{aligned} |([\Xi_\varepsilon(\Theta_\varepsilon^-)] \nabla W_\varepsilon^+(\cdot, t))|_{0, \Omega \setminus B_\rho(t)}^2 &\leq \max_{x \in \Omega \setminus B_{2\rho}(t)} b_\varepsilon(\Theta_\varepsilon^-(x)) |([\Xi_\varepsilon(\Theta_\varepsilon^-)]^{\frac{1}{2}} \nabla W_\varepsilon^+(\cdot, t))|_{0, \Omega \setminus B_\rho(t)}^2 \\ &\leq C(t) b_\varepsilon(1 - 4\rho) \leq C(t) \rho. \end{aligned} \quad (3.25)$$

On noting (3.15) we have for *a.a.* $t \in (0, T)$ that

$$|\Delta^h \Theta_\varepsilon^+(\cdot, t)|_0^2 + \|\underline{U}_\varepsilon^+(\cdot, t)\|_{1,4}^4 \leq C(t). \quad (3.26)$$

It follows from (3.26) and (3.7d), on extracting a further subsequence, that for *a.a.* $t \in (0, T)$ and as $h \rightarrow 0$

$$\Delta^h \Theta_\varepsilon^+(\cdot, t) \rightarrow \Delta \theta(\cdot, t) \quad \text{weakly in } L^2(\Omega), \quad (3.27a)$$

$$\underline{U}_\varepsilon^+(\cdot, t) \rightarrow \underline{u}(\cdot, t) \quad \text{weakly in } \underline{W}^{1,4}(\Omega) \text{ and strongly in } \underline{H}^1(\Omega); \quad (3.27b)$$

see Barrett, Nürnberg, and Styles (2004), as for (3.11a), for details of the former. Combining (2.90), (2.25), (2.4), (3.1b), (3.19) and (3.18) yields for *a.a.* $t \in (0, T)$ and for all $h \leq \widehat{h}_0(\frac{\rho}{2}, t)$ that

$$W_\varepsilon^+(\cdot, t) \equiv -\gamma \Delta^h \Theta_\varepsilon^+(\cdot, t) - \gamma^{-1} \Theta_\varepsilon^-(\cdot, t) + \frac{1}{2} (Q^h[c'(\Theta_\varepsilon^-) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^+) : \underline{\underline{\mathcal{E}}}(U_\varepsilon^+)]) (\cdot, t) \quad \text{on } B_\rho(t). \quad (3.28)$$

If $v_i^{(h)} \in L^4(\Omega)$, $i = 1, 2$, then for any $\eta \in H^2(\Omega)$ we have that

$$\begin{aligned} (Q^h[v_1^h v_2^h] - v_1 v_2, \eta) &= (v_1^h v_2^h - v_1 v_2, \eta) + ((Q^h - I)[v_1^h v_2^h], (I - \pi^h)\eta) \\ &\quad + [(Q^h[v_1^h v_2^h], \pi^h \eta) - (Q^h[v_1^h v_2^h], \pi^h \eta)^h]. \end{aligned} \quad (3.29)$$

It then follows from (3.29), (2.15), (2.18) and a density argument that

$$\begin{aligned} v_i^h &\rightarrow v_i \quad \text{strongly in } L^2(\Omega) \text{ and weakly in } L^4(\Omega), \quad i = 1, 2, \\ \Rightarrow Q^h[v_1^h v_2^h] &\rightarrow v_1 v_2 \quad \text{weakly in } L^2(\Omega). \end{aligned} \quad (3.30)$$

We then have from (3.28), (3.27a,b), (3.16b), (1.4) and (3.30) for *a.a.* $t \in (0, T)$ that as $h \rightarrow 0$

$$W_\varepsilon^+(\cdot, t) \rightarrow -\gamma \Delta \theta(\cdot, t) - \gamma^{-1} \theta(\cdot, t) + \frac{\gamma}{2} [c'(\theta) \mathcal{C} \underline{\underline{\mathcal{E}}}(u) : \underline{\underline{\mathcal{E}}}(u)] (\cdot, t) \quad \text{weakly in } L^2(B_\rho(t)).$$

This together with (3.24) yields

$$W_\varepsilon^+(\cdot, t) \rightarrow w(\cdot, t) \quad \text{weakly in } H^1(B_\rho(t)). \quad (3.31)$$

Combining (3.23), (3.31) and (3.7b) yields for *a.a.* $t \in (0, T)$ that as $h \rightarrow 0$

$$[\Xi_\varepsilon(\Theta_\varepsilon^-) \nabla W_\varepsilon^+(\cdot, t)] \rightarrow b(\theta(\cdot, t)) \nabla w(\cdot, t) \quad \text{weakly in } \underline{L}^2(B_\rho(t)). \quad (3.32)$$

Repeating (3.24) – (3.32) for all $\rho \in (0, \widehat{\rho}_0)$ yields (3.20) and, on noting (3.25) and (3.23), the desired result (3.21). \square

THEOREM. 3.1 *Let the assumptions of Lemma 3.3 hold. Then there exists a subsequence of $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}_h$, where $\{\underline{U}_\varepsilon^+, \Theta_\varepsilon, W_\varepsilon^+\}$ solve $(P_\varepsilon^{h,\tau})$, and functions $\{\underline{u}, \theta, w\}$ satisfying (3.5), (3.10) and (3.20). In addition, as $h \rightarrow 0$ the following hold: (3.6a–c), (3.7a–d), (3.11a–d), (3.16a), (3.16b) for a.a. $t \in (0, T)$, and (3.21). Furthermore, we have that $\{\underline{u}, \theta, w\}$ fulfil $\theta(\cdot, 0) = \theta^0(\cdot)$ in $L^2(\Omega)$ and satisfy (3.8), (3.20) and*

$$\gamma \int_0^T \left\langle \frac{\partial \theta}{\partial t}, \eta \right\rangle dt + \int_{\{|\theta| < 1\}} b(\theta) \nabla w \cdot \nabla \eta \, dx \, dt = 0 \quad \forall \eta \in L^2(0, T; H^1(\Omega)). \quad (3.33)$$

Proof. We need to prove only (3.33). For any $\eta \in H^1(0, T; H^2(\Omega))$ we choose $\chi \equiv \pi^h \eta$ in (3.4b). The desired result (3.33) then follows from (2.18), the embedding $H^1(0, T; X) \hookrightarrow C([0, T]; X)$, (3.12), (2.15), (2.20), (3.6b), (2.78), (3.21) and the denseness of $H^1(0, T; H^2(\Omega))$ in $L^2(0, T; H^1(\Omega))$; see Barrett, Nürnberg, and Styles (2004, Theorem 3.6) for details. \square

4 Numerical Results

Before presenting some numerical results, we briefly state algorithms for solving the resulting system of algebraic equations for $\{\underline{U}_\varepsilon^n, \Theta_\varepsilon^n, W_\varepsilon^n\}$ arising at each time level from the approximation $(P_\varepsilon^{h,\tau})$. As (2.12a) is independent of $\{\Theta_\varepsilon^n, W_\varepsilon^n\}$, we first solve the resulting linear equation to obtain $\underline{U}_\varepsilon^n$. To this end we employ a preconditioned conjugate gradient solver. Then the nonlinear equations (2.12b–c) are solved, using the same ‘‘Gauss–Seidel type’’ iteration as in Barrett, Nürnberg, and Styles (2004, §4).

In order to define the initial shape of the void we introduce the following function. Given $z \in \mathbb{R}^2$ and $R \in \mathbb{R}_{>0}$ we define

$$v(z, R; x) := \begin{cases} -1 & r(x) \leq R - \frac{\gamma\pi}{2} \\ \sin\left(\frac{r(x)-R}{\gamma}\right) & |r(x) - R| < \frac{\gamma\pi}{2}, \\ 1 & r(x) \geq R + \frac{\gamma\pi}{2} \end{cases}, \quad \text{where } r(x) := |x - z|. \quad (4.1)$$

(4.1) represents a circular void with radius R . In line with the asymptotics of the phase field approach, see §1, the interfacial thickness is equal to $\gamma\pi$. For the initial data to (P) we chose θ^0 to be either (i) one circle or (ii) two circles; that is,

$$(i) \quad \theta^0(x) = v(z, R; x) \quad \text{or} \quad (ii) \quad \theta^0(x) = v(z, R; x) + v(\tilde{z}, \tilde{R}; x) - 1. \quad (4.2)$$

We note that in the absence of elastic stresses both these choices of θ^0 are steady states of (P).

Throughout the given domain $\Omega = (-L, L) \times (-L, L)$ is partitioned into right-angled isosceles triangles, such that there are approximately 8 mesh points across the interface. On using the adaptive finite element code Albert 1.0, see Schmidt and Siebert (2001), we implemented the same mesh refinement strategy as in Barrett, Nürnberg, and Styles

(2004). In particular, to improve efficiency we use the approximation $(\tilde{\mathbb{P}}_\varepsilon^{h,\tau})$, see Remark 2.4, and set $\Theta_\varepsilon^0 \equiv \pi^h \theta^0$. Now we have to solve for $\{\Theta_\varepsilon^n, W_\varepsilon^n\}$ only in the interfacial region, $|\Theta_\varepsilon^{n-1}| < 1$. Hence we use a refined mesh with mesh size $h_f = \frac{2^{\frac{3}{2}} L}{N_f}$ in this interfacial region, and a coarser mesh of mesh size $h_c = \frac{2^{\frac{3}{2}} L}{N_c}$ away from the interface. Here N_f and N_c are parameters, see Barrett, Nürnberg, and Styles (2004, §5). Furthermore, we choose N_f such that there are always at least approximately 8 mesh points across the interface in each direction, i.e. $h_f \leq \frac{3\sqrt{2}}{32} \gamma \pi$.

Throughout this section, we restrict ourselves to isotropic elasticity. Hence the assumption (2.34) is satisfied and all our theoretical results in this paper apply. If \mathcal{C} is isotropic, (1.11), then it can be described by its non zero elements

$$\mathcal{C}_{1111} = \mathcal{C}_{2222} = 2\mu + \lambda, \quad \mathcal{C}_{1122} = \lambda, \quad \mathcal{C}_{1212} = \mu; \quad (4.3)$$

where $\mu \in \mathbb{R}_{>0}$ and $\lambda \in \mathbb{R}_{\geq 0}$ are the Lamé moduli.

The following computations are inspired by the results in Bhate, Kumar, and Bower (2000, Figures 9 and 10). They noticed that the void evolution depends strongly on the dimensionless parameter $\Lambda = \frac{S_\infty^2 R}{\beta \varsigma}$ where S_∞ is the maximal stress applied externally, $\beta = \frac{\mu(2\mu+3\lambda)}{\mu+\lambda}$, R is the initial radius of the void, as in (4.1), and ς is surface energy density, which without loss of generality is taken as $\frac{\pi}{2}$ throughout this paper. Unfortunately, the authors did not provide their exact dimensions, but it seems that there $L \approx 4R$ and $R \approx 7\gamma$. Throughout our experiments we set Ω to be the unit square, $L = 0.5$, hence these values correspond to $R \approx \frac{1}{8}$ and $\gamma \approx \frac{1}{56} \approx \frac{1}{18\pi}$. In what follows we set $R = \frac{1}{8}$, $\underline{S} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ for the pure traction boundary condition (1.13) $\Rightarrow S_\infty = 1$, and choose $\gamma \in \{\frac{1}{12\pi}, \frac{1}{24\pi}, \frac{1}{48\pi}\}$. Finally, let \mathcal{C} be defined by (4.3) with $\lambda = \mu = \frac{1}{10\Lambda\pi}$, where $\Lambda \in \{\frac{1}{8}, \frac{1}{5}, \frac{1}{2}\}$.

First, we conducted the following convergence experiments. Setting $c_0(\gamma) = \gamma^2$ in (1.4), we repeated the same experiment with decreasing values of γ , i.e. $\gamma = \frac{1}{12\pi}, \frac{1}{24\pi}, \frac{1}{48\pi}$. In particular, we set $\Lambda = \frac{1}{8}$, $T = 0.02$, $\tau_n = \tau = (\frac{\gamma}{24\pi})^2 \times 10^{-6}$, $\varepsilon = \frac{\gamma}{24\pi} \times 10^{-5}$ and used the appropriate refinement parameters $N_f = \frac{32}{3} \frac{1}{\gamma\pi}$, $N_c = \frac{N_f}{8}$. The steady state solutions for this setup agreed very well for the different values of γ . Hence we are satisfied that the converged solution is very close to the sharp interface limit. See Figure 1, where we superimpose the steady states for $\gamma = \frac{1}{24\pi}$ and $\gamma = \frac{1}{48\pi}$.

For the remaining experiments, we fix $\varepsilon = 10^{-5}$ and set $c_0 = 10^{-3}$ in (1.4). In our first run, we chose $\Lambda = \frac{1}{5}$ as in Bhate, Kumar, and Bower (2000, Fig.9). This yields $\lambda = \mu = \frac{1}{2\pi}$. The other parameters were chosen as follows: $\gamma = \frac{1}{12\pi}$, $T = 0.02$, $\tau_n = \tau = 1.5 \times 10^{-5}$. As initial data we chose (4.2)(i) with $z = (0, 0)$, $R = \frac{1}{8}$. The refinement parameters were $N_f = 128$ and $N_c = 16$. In Figure 2 we plot the zero level sets for $\Theta_\varepsilon(x, t)$ at different times. Note that the last plot is a numerical steady state. Furthermore, the figure contains plots of the principal elastic stress field and the elastic energy at time $t = T$. Here the former is defined as

$$\max\{|\lambda| : \lambda \text{ is an eigenvalue of } c(\Theta_\varepsilon^-) \mathcal{C} \underline{\underline{\mathcal{E}}}(U_\varepsilon^+)\},$$




Figure 1: ($\underline{S} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \Lambda = \frac{1}{8}$) Comparison of zero level sets for $\Theta_\varepsilon(x, t)$ at time $T = 0.02$ for $\gamma = \frac{1}{24\pi}$ and $\gamma = \frac{1}{48\pi}$.

whereas the elastic energy is defined as $c(\Theta_\varepsilon^-) \mathcal{C} \underline{\underline{\xi}}(U_\varepsilon^+) : \underline{\underline{\xi}}(U_\varepsilon^+)$. To simplify matters, both functions were evaluated at the vertices of the triangulation, where we used an arithmetic average of the functions' value on all adjacent triangles. One clearly notices that material is transported away from regions with high elastic energy. To check convergence, we




Figure 2: ($\underline{S} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \Lambda = \frac{1}{5}, \gamma = \frac{1}{12\pi}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0, 1.5 \times 10^{-4}, 1.5 \times 10^{-3}, 0.02$ and elastic stress field and elastic energy at time $t=0.02$.

repeated the same experiment with finer discretization parameters $\tau_n = \tau = 5 \times 10^{-6}$, $N_f = 256$, $N_c = 32$ and the results were graphically indistinguishable from those in Figure 2.

For a smaller interfacial parameter $\gamma = \frac{1}{24\pi}$ we observe a strikingly different behaviour, see Figure 3. The elliptic shape is no longer stable, and this leads to the development of a long slit. Here we see that the condition (3.9) need not always be satisfied in practice. Hence our convergence results for $(P_\varepsilon^{h,\tau})$ and a fixed γ would only hold true, until the void reaches the boundary of the domain and the material is separated into two parts. The evolution in this example indicates that the elastic stresses and the curvature would

become singular in the sharp interface limit. Hence the sharp interface asymptotics, which assumes a bounded curvature, breaks down. These singularities are related to the Asaro–Tiller–Grinfeld instability, see e.g. Asaro and Tiller (1972), Grinfeld (1986) and also Spencer, Voorhees, and Davis (1993). Moreover, it is argued in Kassner, Misbah, Müller, Kappey, and Kohlert (2001) that a phase field model can be interpreted as a regularization of the singularities resulting from these instabilities. In fact they claim that a phase field model might even be more realistic, since it is not clear that the sharp interface model is still plausible in situations where it leads to finite time singularities. We note that our results are in contrast to Bhate, Kumar, and Bower (2000, Fig.9), where the authors used a larger interfacial parameter γ . The discretization parameters used for our computation are $\tau_n = \tau = 2.5 \times 10^{-6}$ and $N_f = 256$, $N_c = 32$.




Figure 3: ($\underline{S} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $\Lambda = \frac{1}{5}$, $\gamma = \frac{1}{24\pi}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0$, 1.5×10^{-4} , 1.5×10^{-3} , 3×10^{-3} , 3.75×10^{-3} , 5.25×10^{-3} and elastic stress field and elastic energy at time $t=3.75 \times 10^{-3}$.

The next run is for $\Lambda = \frac{1}{2}$ as in Bhate, Kumar, and Bower (2000, Fig.10), i.e. $\lambda = \mu = \frac{1}{5\pi}$. A computation for $\gamma = \frac{1}{12\pi}$, $T = 4 \times 10^{-5}$, $\tau_n = \tau = 5 \times 10^{-7}$ and refinement parameters $N_f = 128$, $N_c = 16$ can be seen in Figure 4. Again we can observe a slightly different evolution for a smaller value of γ , see Figure 5. In particular, the developing cusps appear sharper and less smoothed out. One can again clearly see that material is transported away from regions with high elastic energy. The parameters for this computation were $\gamma = \frac{1}{24\pi}$, $T = 10^{-5}$, $\tau_n = \tau = 10^{-7}$, $N_f = 256$ and $N_c = 32$.

A run with parameters as in Figure 2 but $\underline{S} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ can be seen in Figure 6, where the last plot is a numerical steady state. If we choose a smaller interfacial parameter $\gamma = \frac{1}{24\pi}$, the elastic effect tends to be more pronounced and the steady state shape is slightly different, see Figure 7, where we used the same discretization parameters as in Figure 3. The last plot is a numerical steady state and it is noteworthy that the steady




Figure 4: ($\underline{\underline{S}} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $\Lambda = \frac{1}{2}$, $\gamma = \frac{1}{12\pi}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0, 3 \times 10^{-6}, 10^{-5}, 4 \times 10^{-5}$ and elastic stress field and elastic energy at time $t = 4 \times 10^{-5}$.

state is nonconvex in contrast to Wulff shapes which are minimizers of an anisotropic surface energy under a volume constraint.

For our last example, we chose \mathcal{C} such that $\mathcal{C} \underline{\underline{\mathcal{E}}}(\underline{u}) = \underline{\underline{\mathcal{E}}}(\underline{u})$, i.e. $\mu = \frac{1}{2}$ and $\lambda = 0$, and set $\underline{\underline{S}} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. Starting with two initially circular voids, the presence of elastic stresses leads to a vertical split in the material, see Figure 8. We used the following parameters for $(\tilde{\mathbf{P}}_\varepsilon^{h,\tau})$: $\gamma = \frac{1}{24\pi}$, $T = 5 \times 10^{-5}$ and $\tau_n = \tau = 10^{-7}$. As initial data we chose (4.2)(ii) with $z = -\tilde{z} = (0, 0.23)$, $R = \tilde{R} = 0.18$. The refinement parameters were $N_f = 256$ and $N_c = 32$.

Further results, including simulations modelling the (combined) effect of surface diffusion, an electric field, grain boundaries and anisotropic elasticity will be reported on elsewhere, see Barrett, Garcke, and Nürnberg (2004b), where we also discuss applications to epitaxial growth.

Acknowledgement Part of this work was carried out while the authors participated in the 2003 programme *Computational Challenges in Partial Differential Equations* at the Isaac Newton Institute, Cambridge, UK. This work was also supported by the RTN Programme Fronts-Singularities, HPRN-CT-2002-00274 and the DFG Priority Program Analysis, Modeling and Simulation of Multiscale Problems.

References

- Adams, R. A. and J. Fournier (1977). Cone conditions and properties of Sobolev spaces. *J. Math. Anal. Appl.* 61, 713–734.
- Asaro, R. J. and W. A. Tiller (1972). Interface morphology development during stress




Figure 5: ($\underline{S} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $\Lambda = \frac{1}{2}$, $\gamma = \frac{1}{24\pi}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0, 3 \times 10^{-6}, 5 \times 10^{-5}, 10^{-5}$ and elastic stress field and elastic energy at time $t=10^{-5}$.

corrosion cracking: Part i. via surface diffusion. *Metall. Trans. 3*, 1789–1796.

Barrett, J. W. and J. F. Blowey (2001). Finite element approximation of a degenerate Allen–Cahn/Cahn–Hilliard system. *SIAM J. Numer. Anal.* *39*, 1598–1624.

Barrett, J. W., J. F. Blowey, and H. Garcke (1998). Finite element approximation of a fourth order nonlinear degenerate parabolic equation. *Numer. Math.* *80*, 525–556.

Barrett, J. W., J. F. Blowey, and H. Garcke (1999). Finite element approximation of the Cahn–Hilliard equation with degenerate mobility. *SIAM J. Numer. Anal.* *37*, 286–318.

Barrett, J. W., J. F. Blowey, and H. Garcke (2001). On fully practical finite element approximations of degenerate Cahn–Hilliard systems. *M2AN Math. Model. Numer. Anal.* *35*, 713–748.

Barrett, J. W., H. Garcke, and R. Nürnberg (2003). Finite element approximation of surfactant spreading on a thin film. *SIAM J. Numer. Anal.* *41*, 1427–1464.

Barrett, J. W., H. Garcke, and R. Nürnberg (2004a). A phase field model for electromigration of intergranular voids. (in preparation).

Barrett, J. W., H. Garcke, and R. Nürnberg (2004b). Phase field models for stress and electromigration induced surface diffusion with applications to epitaxial growth and void evolution. (in preparation).

Barrett, J. W., S. Langdon, and R. Nürnberg (2004). Finite element approximation of a sixth order nonlinear degenerate parabolic equation. *Numer. Math.* *96*, 401–434.

Barrett, J. W. and R. Nürnberg (2004). Convergence of a finite element approximation of surfactant spreading on a thin film in the presence of van der Waals forces. *IMA J. Numer. Anal.* *24*, 323–363.




Figure 6: ($\underline{S} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\Lambda = \frac{1}{5}$, $\gamma = \frac{1}{12\pi}$) Zero level sets for $\Theta_\epsilon(x, t)$ at times $t = 0, 1.5 \times 10^{-4}, 1.5 \times 10^{-3}, 0.02$ and elastic stress field and elastic energy at time $t=0.02$.

- Barrett, J. W., R. Nürnberg, and V. Styles (2004). Finite element approximation of a void electromigration model. *SIAM J. Numer. Anal.* (to appear).
- Bergh, J. and J. Löfström (1976). *Interpolation Spaces, an Introduction*. Berlin: Springer.
- Bhate, D. N., A. Kumar, and A. F. Bower (2000). Diffuse interface model for electromigration and stress voiding. *J. Appl. Phys.* *87*, 1712–1721.
- Blowey, J. F. and C. M. Elliott (1991). The Cahn–Hilliard gradient theory for phase separation with non-smooth free energy. Part I: Mathematical analysis. *European J. Appl. Math.* *2*, 233–279.
- Bower, A. and D. Craft (1998). Analysis of failure mechanisms in the interconnect lines of microelectronic circuits. *Fat. Frac. Eng. Mat. Struct.* *21*, 611–630.
- Brenner, S. C. and L. R. Scott (2002). *The Mathematical Theory of Finite Element Methods (second edition)*. New York: Springer.
- Cahn, J. W., C. M. Elliott, and A. Novick-Cohen (1996). The Cahn–Hilliard equation with a concentration dependent mobility: motion by minus the Laplacian of the mean curvature. *European J. Appl. Math.* *7*, 287–301.
- Carrive, M., A. Miranville, and A. Piétrus (2000). The Cahn–Hilliard equation for deformable elastic media. *Adv. Math. Sci. Appl.* *10*, 539–569.
- Ciarlet, P. G. (1988). *Numerical Linear Algebra and Optimisation*. Cambridge: Cambridge University Press.
- Costabel, M., M. Dauge, and Y. Lafranche (2001). Fast semi-analytic computation of elastic edge singularities. *Comp. Meth. Appl. Mech. Eng.* *190*, 2111–2134.
- Elliott, C. M. (1997). Approximation of curvature dependent interface motion. In I. S. Duff and G. A. Watson (Eds.), *The State of the Art in Numerical Analysis*, New




Figure 7: ($\underline{S} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\Lambda = \frac{1}{5}$, $\gamma = \frac{1}{24\pi}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0, 1.5 \times 10^{-4}, 1.5 \times 10^{-3}, 0.02$ and elastic stress field and elastic energy at time $t=0.02$.

York, pp. 407–440. Oxford University Press.

Elliott, C. M. and H. Garcke (1996). On the Cahn–Hilliard equation with degenerate mobility. *SIAM J. Math. Anal.* *27*, 404–423.

Elliott, C. M. and H. Garcke (1997). Existence results for diffusive surface motion laws. *Adv. Math. Sci. Appl.* *7*, 465–488.

Escher, J., U. F. Mayer, and G. Simonett (1998). The surface diffusion flow for immersed hypersurfaces. *SIAM J. Math. Anal.* *29*, 1419–1433.

Fried, E. and M. Gurtin (1994). Dynamic solid-solid transitions with phase characterized by an order parameter. *Physica D* *72*, 287–308.

Garcke, H. (2000). On mathematical models for phase separation in elastically stressed solids. Habilitation, University Bonn.

Garcke, H. (2003). On Cahn–Hilliard systems with elasticity. *Proc. Roy. Soc. Edinburgh* *133 A*, 307–331.

Garcke, H. (2004). On a Cahn–Hilliard model for phase separation with elastic misfit. (submitted for publication).

Garcke, H., M. Rumpf, and U. Weikard (2001). The Cahn–Hilliard equation with elasticity — finite element approximation and qualitative studies. *Interfaces Free Bound.* *3*, 101–118.

Garcke, H. and U. Weikard (2004). Numerical approximation of the Cahn–Larché equation. (submitted).

Giaquinta, M. and G. Modica (1979). Regularity results for some classes of higher order nonlinear elliptic systems. *J. Reine Angew. Math.* *311–312*, 145–169.

Grinfeld, M. A. (1986). Instability of the separation boundary between a non-hydrostatically stressed elastic body and a melt. *Soviet Phys. Dokl.* *31*, 831–834.




Figure 8: ($\underline{S} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$) Zero level sets for $\Theta_\varepsilon(x, t)$ at times $t = 0, 10^{-5}, 2 \times 10^{-5}, 5 \times 10^{-5}$ and elastic stress field and elastic energy at time $t=10^{-5}$.

- Grisvard, P. (1989). Singularités en élasticité. *Arch. Ration. Mech. Anal.* *107*, 157–180.
- Grün, G. (2003). On the convergence of entropy consistent schemes for lubrication type equations in multiple space dimensions. *Math. Comp.* *72*, 1251–1279.
- Grün, G. and M. Rumpf (2000). Nonnegativity preserving numerical schemes for the thin film equation. *Numer. Math.* *87*, 113–152.
- Gungor, M. R., D. Maroudas, and L. J. Gray (1998). Effects of mechanical stress on electromigration-driven transgranular void dynamics in passivated metallic thin films. *Appl. Phys. Lett.* *73*, 3848–3850.
- Gurtin, M. E. (1972). The linear theory of elasticity. In S. Flügge and C. Truesdell (Eds.), *Handbuch der Physik, Vol. VIa/2*. Berlin: Springer.
- Kassner, K., C. Misbah, J. Müller, J. Kappey, and P. Kohlert (2001). Phase-field modeling of stress-induced instabilities. *Phys. Rev. E* *63*, 036117 (27 pages).
- Leo, P. H., J. S. Lowengrub, and H. J. Jou (1998). A diffuse interface model for microstructural evolution in elastically stressed solids. *Acta Mater.* *46*, 2113–2130.
- Mosolov, P. P. and V. P. Mjasnikov (1971). A proof of Korn’s inequality. *Soviet Math. Dokl.* *12*, 1618–1622.
- Nečas, J. and I. Hlaváček (1981). *Mathematical Theory of Elastic and Elasto-Plastic Bodies: An Introduction*. Amsterdam: Elsevier.
- Rannacher, R. and R. Scott (1982). Some optimal error estimates for piecewise linear finite element approximations. *Math. Comp.* *38*, 437–445.
- Schmidt, A. and K. G. Siebert (2001). Albert — software for scientific computations and applications. *Acta Math. Univ. Comenian.* *70*, 105–122.
- Seif, J. B. (1973). On the Green’s function for the biharmonic equation in an infinite wedge. *Trans. Amer. Math. Soc.* *182*, 241–260.

- Simon, J. (1987). Compact sets in the space $L^p(0, T; B)$. *Ann. Math. Pura. Appl.* 146, 65–96.
- Spencer, B. J., P. W. Voorhees, and S. H. Davis (1993). Morphological instability in epitaxially strained dislocation-free solid films: Linear stability theory. *J. Appl. Phys.* 73, 4955–4970.
- Xia, L., A. F. Bower, Z. Suo, and C. Shih (1997). A finite element analysis of the motion and evolution of voids due to strain and electromigration induced surface diffusion. *J. Mech. Phys. Solids* 45, 1473–1493.
- Zhornitskaya, L. and A. L. Bertozzi (2000). Positivity preserving numerical schemes for lubrication-type equations. *SIAM J. Numer. Anal.* 37, 523–555.